Clinical Trial Protocol
Iranian Registry of Clinical Trials

08 Nov 2026

The study of the Effect of acupressure on Hugo point on delivery pain of multiparous women

Protocol summary

Summary

This study is a clinical trial which will be done in labor ward of Abadan 17 Shahrivar hospital. The study objective is to investigate the effect of acupressure on Hugo point on delivery pain of multiparous women. Study population includes the multiparous women who will give birth vaginally at 17 Shahrivar hospital with a moderate to severe delivery pain in 2 hours after delivery (based on the standard visual pain scale assessment). Inclusion criteria include: Vaginal delivery; second to fourth delivery; gestational age between 37 and 42 weeks; complaint of moderate to severe delivery pain; infant birth weight between 2500 to 4000 gram and body mass index between 18.5 and 29. Exclusion criteria include: Hard and prolonged delivery; addiction; history of cesarean section and intra-abdominal surgery and history of postpartum hemorrhage. Sample size has been calculated as 62 women (31 women in each group). Sampling will be done using blocked randomization sampling with blocks' size of 6. SPPS software (random number generation) will be used for randomization. Also, this study is a single blind trial in which the participants will not inform from their group. To conduct the study, first the delivery pain of women who meet the eligibility criteria of study and have the willingness to participate in the study will be assessed at 2 hours after delivery and those with moderate to severe pain will be randomly allocated to intervention or control group. For intervention group, the delivery pain will be assessed using standard visual pain scale assessment and recorded. Also, pain duration of women will be asked from them and recorded. Then, Hugo point will be pressured by researcher for 4 cycles. Each pressure course will take 20 minutes in which, first the Hugo point will be pressured lightly and the pressure will be increased gradually in such a way that the pressure reaches its maximum within 30 seconds. This maximum pressure will be kept for one minute and then will be reduced so that the pressure will be removed from Hugo point within 30 seconds. This pressure will be repeated for 20 minutes (10 times). Then, the delivery pain again will be assessed and recorded 100 minutes after the termination of the each pressure course, the pain duration also will be asked and recorded in that time and a new pressure course will be started. The pressure courses will be continued up to 8 hours after delivery. In control group, a similar intervention will be done with the exception that an irrelevant point will be pressured rather than Hugo point. Finally, the pain severity and duration of control group, before and after intervention will be compared with those of control group.

General information

Acronym

IRCT registration information

IRCT registration number: **IRCT2016112321670N2**
Registration date: **2017-01-31, 1395/11/12**
Registration timing: **registered_while_recruiting**

Last update:
Update count: **0**

Registration date

2017-01-31, 1395/11/12

Registrant information

Name
Elham Bahrami
Name of organization / entity
Shahid Sadoughi University of Medical Sciences
Country
Iran (Islamic Republic of)
Phone
+98 35 3822 4000
Email address
bahrami_hmrc@ssu.ac.ir

Recruitment status

Recruitment complete

Funding source

Investigator

Expected recruitment start date
2017-01-19, 1395/10/30
Expected recruitment end date
2017-07-21, 1396/04/30
Actual recruitment start date
empty
Actual recruitment end date
empty
Trial completion date
empty

Scientific title
The study of the Effect of acupressure on Hugo point on delivery pain of multiparous women

Public title
The study of the effect of acupressure on Hugo point on delivery pain of multiparous women
Purpose
Supportive
Inclusion/Exclusion criteria
Inclusion criteria: Vaginal delivery; second to fourth pregnancy; gestational age between 37 to 42 weeks; complaint of moderate to severe delivery pain based on the standard visual pain scale assessment; infant birth weight between 2500 to 4000 gram; body mass index between 18.5 to 29 Exclusion criteria: Hard and prolonged delivery; addiction; history of cesarean section and intra-abdominal surgery; history of postpartum hemorrhage
Age
No age limit
Gender
Female

Phase
N/A
Groups that have been masked
No information
Sample size
Target sample size: **62**
Randomization (investigator's opinion)
Randomized
Randomization description
Blinding (investigator's opinion)
Single blinded
Blinding description
Placebo
Not used
Assignment
Single
Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee
Name of ethics committee
Ahvaz Jondi Shapoor University of Medical Sciences
Street address
Ahvaz Jondi Shapoor University of Medical Sciences, University town, Ahvaz, Iran
City
Ahvaz
Postal code
6135715794
Approval date
2016-10-22, 1395/08/01
Ethics committee reference number
IR.AJUMS.REC.1395.432

Health conditions studied

1

Description of health condition studied
The relationship of acupressure on Hugo point and delivery pain
ICD-10 code
047.1
ICD-10 code description
False labour at or after 37 completed weeks of gestation

Primary outcomes

1

Description
The primary outcome measure of this study is severity of delivery pain.
Timepoint
The severity of delivery pain will be assessed before and after each pressure course.
Method of measurement
Severity of delivery pain will be assessed using standard visual pain scale assessment.

Secondary outcomes

1

Description
Secondary outcome measure of this study is the duration of delivery pain.
Timepoint
Duration of delivery pain will be assessed before and after of each pressure course.
Method of measurement
Duration of delivery pain will be asked from participants and recorded.

Intervention groups

1

Description
Intervention group: For this group, Hugo point will be pressured by researcher for 4 cycles. Each pressure course will take 20 minutes in which, first the Hugo point will be pressured lightly and the pressure will be increased gradually in such a way that the pressure reaches its maximum within 30 seconds. This maximum pressure will be kept for one minute and then will be reduced so that the pressure will be removed from Hugo point within 30 seconds. This pressure will be repeated for 20 minutes (10 times). Then, the delivery pain again will be assessed and recorded 100 minutes after the termination of the each pressure course, the pain duration also will be asked and recorded in that time and a new pressure course will be started. The pressure courses will be continued up to 8 hours after delivery.

Category
Other

2

Description
Control group: In this group, an irrelevant point other than Hugo point will be pressured by researcher for 4 cycles. Each pressure course will take 20 minutes in which, first the Hugo point will be pressured lightly and the pressure will be increased gradually in such a way that the pressure reaches its maximum within 30 seconds. This maximum pressure will be kept for one minute and then will be reduced so that the pressure will be removed from under pressure point within 30 seconds. This pressure will be repeated for 20 minutes (10 times). Then, the delivery pain again will be assessed and recorded 100 minutes after the termination of the each pressure course, the pain duration also will be asked and recorded in that time and a new pressure course will be started. The pressure courses will be continued up to 8 hours after delivery.

Category
Other

Recruitment centers

1

Recruitment center
Name of recruitment center
Abadan 17 Shahrivar hospital
Full name of responsible person
Shahla Afravi (Master of Science, midwifery)
Street address
17 Shahrivar hospital, Shatit, Abadan, Khoozestan, Iran
City
Abadan

Sponsors / Funding sources

1

Sponsor
Name of organization / entity
Abadan 17 Shahrivar hospital
Full name of responsible person
Shahla Afravi
Street address
17 Shahrivar hospital, Shatit, Abadan, Khoozestan, Iran
City
Abadan
Grant name
Grant code / Reference number
Is the source of funding the same sponsor organization/entity?
Yes
Title of funding source
Abadan 17 Shahrivar hospital
Proportion provided by this source
100
Public or private sector
empty
Domestic or foreign origin
empty
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
empty

Person responsible for general inquiries

Contact
Name of organization / entity
Abadan 17 Shahrivar hospital
Full name of responsible person
Shahla Afravi
Position
Master of Science, midwifery
Other areas of specialty/work
Street address
17 Sharivar hospital, Shatit, Abadan, Khoozestan, Iran
City
Abadan
Postal code
0631891331
Phone
+98 61 5335 4546
Fax
+98 61 5335 4540
Email
af.shahla@yahoo.com
Web page address

Person responsible for scientific inquiries

Contact
Name of organization / entity
Abadan 17 Shahrivar hospital
Full name of responsible person

Shahla Afravi

Position

Master of Science, midwifery

Other areas of specialty/work

Street address

17 Shahrivar hospital, Shatit, Abadan, Khoozestan, Iran

City

Abadan

Postal code

0631891331

Phone

+98 61 5335 4546

Fax

+98 61 5335 4540

Email

af.shahla@yahoo.com

Web page address

Person responsible for updating data

Contact

Name of organization / entity

Abadan 17 Shahrivar hospital

Full name of responsible person

Shahla Afravi

Position

Master of Science, midwifery

Other areas of specialty/work

Street address

17 Shahrivar hospital, Shatit, Abadan, Khoozestan, Iran

City

Abadan

Postal code

0631891331

Phone

+98 61 5335 4546

Fax

+98 61 5335 4540

Email

af.shahla@yahoo.com

Web page address

Sharing plan

Deidentified Individual Participant Data Set (IPD)

empty

Study Protocol

empty

Statistical Analysis Plan

empty

Informed Consent Form

empty

Clinical Study Report

empty

Analytic Code

empty

Data Dictionary

empty