

Clinical Trial Protocol

Iranian Registry of Clinical Trials

02 Jun 2026

Comparison the prevalence of sore throat due to endotracheal intubation with videolaryngoscope (Glidoscope) and conventional laryngoscope Macintosh and placement the LMA in elective surgeries

Protocol summary

Study aim

Comparison the prevalence of sore throat due to endotracheal intubation with videolaryngoscope (Glidoscope) and conventional laryngoscope Macintosh and placement the LMA in elective surgeries

Design

Study design: clinical trial. Number of groups: 3 Total sample size: 90 patients. Sample size in each group: 30 people Randomization: random packs that opened precisely before the start of surgery. Blinding: a person outside the study and unaware of the study process (double-blind study).

Settings and conduct

Location: Shahid Rajaei & Velayat hospitals in Qazvin city, Iran. Study Methods: patients were randomly assigned to one of the study groups. Patient's monitoring was performed using ECG (electrocardiogram), pulse oximetry and barometric measurements. All patients received 2 mg/kg body weight fentanyl and 0.2 mg/kg body weight midazolam as a precursor and then propofol was given at a dose of 2.5-2 mg / kg body weight and atracurium at a dose of 0.5 mg/kg of body weight, anesthetic induction was prescribed. Patients were treated with 100% oxygenation mask for 3 minutes; then the G group was subjected to intubation with blade number 3; the M group was also implanted with blade 3 of the Macintosh laryngoscope and the L group was placed under the LMA placement. After fixing the tube or LMA, an oxygen concentration of 100% plus TIVA (propofol infusion) was administered to each of the three groups of patients for anesthesia (N2O not prescribed for any patient), and repeated fentanyl injection was used for maintaining analgesia. The number of patients with sore throat in the 8-6 hours and 24 hours after surgery was recorded in all three groups using questionnaire. The length of the laryngoscopy (that is measured by the number of seconds from the time the laryngoscope

enters the mouth, until the tracheal tube passes) was recorded via vocal cords. For each patient, the duration of the laryngoscopy until the insertion of the tracheal tube and the number of attempts to intubate by seconds were measured and recorded. The length of the LMA placement was measured and recorded from the time it was inserted into the mouth until it was placed in Pharynx. Also in all patients, hemodynamic parameters including heart rate, systolic blood pressure, diastolic blood pressure and mean blood pressure were recorded one minute before induction of anesthesia and also in the 3rd and 5th minutes after intubation or LMA insertion.

Participants/Inclusion and exclusion criteria

Inclusion criteria: presence of ASA (American Society of Anesthesiology Status) Class I or II; age between 20 to 40 years old; patients undergoing any elective surgery (except for head and neck surgery) and requiring general anesthesia. Non-admission criteria: aspiration and vomiting during induction of anesthesia and laryngoscopy; the presence of any lesion or active disease in the head and neck area and the airway such as pharyngitis or laryngeal mass during pre-anesthetic visit; head and neck surgery; cardiovascular, respiratory, neovascular and cardiovascular and rheumatic disorders; high risk of aspirating (diabetic patients, incomplete NPO time, addicted patients); history of difficult Intubation Length of surgery for more than 2 hours; possible problems for intubation in the clinical examination; length of intubation or LMA insertion time greater than 20 seconds.

Intervention groups

1) Direct laryngoscopy group via Macintosh blade (M) 2) Indirect laryngoscopy group via glidoscope (G) 3) Group (L) by inserting LMA

Main outcome variables

Sore throat at intervals of 8-6 hours and 24 hours after intubation

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20180207038661N1**

Registration date: **2018-02-18, 1396/11/29**

Registration timing: **retrospective**

Last update: **2018-02-18, 1396/11/29**

Update count: **0**

Registration date

2018-02-18, 1396/11/29

Registrant information

Name

Mohammad Ali Masoumifar

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 28 3332 0107

Email address

mebtehaj@qums.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2018-01-21, 1396/11/01

Expected recruitment end date

2018-03-11, 1396/12/20

Actual recruitment start date

2017-06-23, 1396/04/02

Actual recruitment end date

2017-12-15, 1396/09/24

Trial completion date

empty

Scientific title

Comparison the prevalence of sore throat due to endotracheal intubation with videolaryngoscope (Glidoscope) and conventional laryngoscope Macintosh and placement the LMA in elective surgeries

Public title

Comparison sore throats caused by the three main techniques of intracranial intubation

Purpose

Treatment

Inclusion/Exclusion criteria

Inclusion criteria:

Presence of ASA (American Society of Anesthesiology Status) Class I or II Age between 20 to 40 years old Patients undergoing any elective surgery (except for head and neck surgery) and requiring general anesthesia.

Exclusion criteria:

Aspiration and vomiting during induction of anesthesia and laryngoscopy The presence of any lesion or active disease in the head and neck area and the airway such as pharyngitis or laryngeal mass during pre-anesthetic

visit Head and neck surgery Cardiovascular, respiratory, neovascular and cardiovascular and rheumatic disorders High risk of aspirating (diabetic patients, incomplete NPO time, addicted patients) History of difficult Intubation Length of surgery for more than 2 hours Possible problems for intubation in the clinical examination Length of intubation or LMA insertion time greater than 20 seconds

Age

From **20 years** old to **40 years** old

Gender

Both

Phase

N/A

Groups that have been masked

- Participant
- Investigator
- Outcome assessor

Sample size

Target sample size: **90**

Actual sample size reached: **90**

Randomization (investigator's opinion)

Randomized

Randomization description

In order to randomization and placement of patients in one of the study groups we used random packets. In this way, the researcher randomly opened one of the envelopes before the anesthetic induction and at that moment the technique was chosen for the patient.

Blinding (investigator's opinion)

Double blinded

Blinding description

Packets used for randomization, were prepared by someone else who were not included in the study.

Placebo

Not used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics Committee in Biomedical Research of Qazvin University of Medical Sciences

Street address

Qazvin University of Medical Sciences, Bahonar Bulivard, Qazvin, Iran

City

Qazvin

Province

Qazvin

Postal code

34197-59811

Approval date

2016-11-26, 1395/09/06

Ethics committee reference number

IR.QUMS.REC.1395.192

Health conditions studied

1

Description of health condition studied

Patients undergoing elective surgeries (Non-emergent)

ICD-10 code

Z41.9

ICD-10 code description

Procedure for purposes other than remedying health state, unspecified

Primary outcomes

1

Description

Sore-throat

Timepoint

6-8 hours and 24 hours after intubation

Method of measurement

Asking question from patient

Secondary outcomes

1

Description

Systolic Blood Pressure

Timepoint

Before starting, Min 3 after starting, Min5 after starting

Method of measurement

Barometer

2

Description

Diastolic Blood Pressure

Timepoint

Before starting, Min 3 after starting, Min5 after starting

Method of measurement

Barometer

3

Description

Mean Blood Pressure

Timepoint

Before starting, Min 3 after starting, Min5 after starting

Method of measurement

Mathematical Calculation

4

Description

Heart Rate

Timepoint

Before starting, Min 3 after starting, Min5 after starting

Method of measurement

Heart Monitoring

5

Description

Duration of intubation or LMA embedding

Timepoint

Once, During the procedure

Method of measurement

Chronometer

6

Description

The duration between laryngoscopy until the insertion of the tracheal tube

Timepoint

Once, During the procedure

Method of measurement

Chronometer

7

Description

The number of attempts for intubation or LMA placement

Timepoint

Once, During the procedure

Method of measurement

Anesthesiologist

Intervention groups

1

Description

Group 1: Direct laryngoscopy with Macintosh blade

Category

Treatment - Surgery

2

Description

Group 2: Indirect laryngoscopy by glidoscope

Category

Treatment - Surgery

3

Description

Group 3: Laryngeal mask airway (LMA) insertion

Category

Treatment - Surgery

Recruitment centers

1

Recruitment center

Name of recruitment center

Shahid Rajae Hospital

Full name of responsible person

Mohammad Ali Masoumi far

Street address

Shahid Rajaei Hospital, Shahid Gharani Street, End of Gargan Street, Qazvin

City

Qazvin

Province

Qazvin

Postal code

3419759811

Phone

+98 28 3332 0105

Email

hosrajaee@qums.ac.ir

Web page address

<http://hosrajaee.qums.ac.ir/Portal/Home/>

Grant name**Grant code / Reference number****Is the source of funding the same sponsor organization/entity?**

Yes

Title of funding source

Qazvin University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin**Type of organization providing the funding**

Academic

2**Recruitment center****Name of recruitment center**

Velayat Hospital

Full name of responsible person

Mohammad Ali Masoumi Far

Street address

Velayat Hospital, 22 Bahman Bulivard, Elahieh, Minoodar, Qazvin

City

Qazvin

Province

Qazvin

Postal code

3419759811

Phone

+98 28 3379 0620

Email

Info@hosvelayat.qums.ac.ir

Web page address

<http://hosvelayat.qums.ac.ir/Portal/Home/Default.aspx>

Person responsible for general inquiries**Contact****Name of organization / entity**

Qazvin University of Medical Sciences

Full name of responsible person

Mohammad Ali Masoumi far

Position

دستیار بیهوشی

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Street address

Qazvin University of Medical Sciences, Shahid Bahonar Blvd. , Qazvin

City

Qazvin

Province

Qazvin

Postal code

3419759811

Phone

+98 28 3333 6001

Email

ma.masoumifar@gmail.com

Sponsors / Funding sources**1****Sponsor****Name of organization / entity**

Qazvin University of Medical Sciences

Full name of responsible person

Dr. Amir Peymani

Street address

Mavaddat Street, Shahid Beheshti Blvd, Qazvin

City

Qazvin

Province

Qazvin

Postal code

1391134156

Phone

+98 28 3333 7006

Email

research.dpt@qums.ac.ir

Person responsible for scientific inquiries**Contact****Name of organization / entity**

Qazvin University of Medical Sciences

Full name of responsible person

Mohammad Ali Masoumi far

Position

Anesthesiologist Assistant

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Street address

Qazvin University of Medical Sciences, Shahid

Bahonar Blvd. , Qazvin

City

Qazvin

Province

Qazvin

Postal code

3419759811

Phone

+98 28 3333 6001

Email

ma.masoumifar@gmail.com

Person responsible for updating data

Contact

Name of organization / entity

Qazvin University of Medical Sciences

Full name of responsible person

Mohammad Ali Masoumi far

Position

Anesthesiologist Assistant

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Street address

Qazvin University of Medical Sciences, Shahid
Bahonar Blvd. , Qazvin

City

Qazvin

Province

Qazvin

Postal code

3419759811

Phone

+98 28 3333 6001

Email

ma.masoumifar@gmail.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

No - There is not a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Yes - There is a plan to make this available

Analytic Code

No - There is not a plan to make this available

Data Dictionary

No - There is not a plan to make this available

Title and more details about the data/document

I have not yet decided on this.

When the data will become available and for how long

I have not yet decided on this.

To whom data/document is available

I have not yet decided on this.

Under which criteria data/document could be used

I have not yet decided on this.

From where data/document is obtainable

I have not yet decided on this.

What processes are involved for a request to access data/document

I have not yet decided on this.

Comments