

Clinical Trial Protocol

Iranian Registry of Clinical Trials

28 Jun 2026

Comparison of newly Presented Suturing technique with Conventional Suturing on Severity of Complications Following Mandibular fully Impacted third molar Surgery

Protocol summary

Study aim

The innovative suture technique with conventional suture during mandibular fully impacted third molar surgery on the severity of postoperative complications will compare

Design

Controlled clinical trial with a parallel group, double blinded, randomized with statistical software using blocks of six

Settings and conduct

This study carried out in department of Surgery, Faculty of Dentistry, Semnan University of Medical Sciences The suture pattern will be selected randomly in the first surgery and will not be shown to the surgeon prior to the start of suturing and finishing the surgical procedure. The patient and measurer will be blind about the type of suture

Participants/Inclusion and exclusion criteria

Inclusion criteria: Patients aged between 20 and 30 Both-sided and symmetrical impaction of mandibular third molar tooth that are classified in the same class by Pell & Gregory Meeting ASA1 systemic conditions Collaborative patients with frequent visits for follow-up Patients without periodontal disease The exclusion criteria of the study include: Pathologic lesion in the impacted tooth Difference in follicle size around the crown of each person's sides Taking steroids, drugs, alcohol, contraceptives Pregnancy and lactation History of delayed wound healing Failure to complete follow-up sessions

Intervention groups

In the control side simple loop suture. In the case side, first the thread will be twisted around the needle holder in clockwise direction A) Then, with the aid of the needle holder the middle of the loose thread is taken and passed around the needle holder through the twisted thread to form a loop B) The needle holder is passed

through the loop into the free end of the thread and pulled to complete the knot

Main outcome variables

Pain: maximal: oral opening rate: facial swelling: the 7th tooth distal pocket depth: clinical attachment loss in 7th tooth distal: dry socket: dehiscence

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20181021041404N1**

Registration date: **2018-11-14, 1397/08/23**

Registration timing: **registered_while_recruiting**

Last update: **2018-11-14, 1397/08/23**

Update count: **0**

Registration date

2018-11-14, 1397/08/23

Registrant information

Name

MohammadHosein Mansourian

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 8821 4780

Email address

mh.mansourian95@gmail.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2018-11-07, 1397/08/16

Expected recruitment end date

2019-01-05, 1397/10/15

Actual recruitment start date
empty

Actual recruitment end date
empty

Trial completion date
empty

Scientific title
Comparison of newly Presented Suturing technique with Conventional Suturing on Severity of Complications Following Mandibular fully Impacted third molar Surgery

Public title
Comparison of Innovative Suture with Conventional Suture on Severity of Complications Following Mandibular fully Impacted third molar Surgery

Purpose
Treatment

Inclusion/Exclusion criteria
Inclusion criteria:
Both-sided and symmetrical impaction of mandibular third molar tooth that are classified in the same class by Pell & Gregory Meeting ASA1 systemic conditions Non-smokers patients Collaborative patients with frequent visits for follow up Patients without periodontal disease patients with proper oral hygiene patients aged between 20 and 30
Exclusion criteria:
Pathologic lesion in the impacted tooth Difference in follicle size around the crown of each person's sides Surgical time exceeding 25 minutes More than 5 minutes of surgical difference on both sides Taking steroids, drugs, alcohol, contraceptives Pregnancy and lactation History of delayed wound healing Failure to complete follow up sessions

Age
From **20 years** old to **30 years** old

Gender
Both

Phase
N/A

Groups that have been masked

- Participant
- Care provider
- Outcome assessor
- Data analyser
- Data and Safety Monitoring Board

Sample size
Target sample size: **40**
More than 1 sample in each individual
Number of samples in each individual: **2**
Right and left mandibular fully impacted third molar

Randomization (investigator's opinion)
Randomized

Randomization description
In order to randomize, patients are divided into two groups using the blocks of six that made by statistical software: In the first group, the first surgery is on the lower right jaw, and in the second group, the first surgery is performed on the lower left jaw. All surgeries will be

performed by an oral and maxillofacial surgeon and in one room with an assistant. The suture pattern will be selected randomly in the first surgery and will not be shown to the surgeon prior to the start of suturing and finishing the surgical procedure.

Blinding (investigator's opinion)

Double blinded

Blinding description

The suture pattern will not be shown to the surgeon(main researcher) prior to the start of suturing and finishing the surgical procedure. The patient and measurer will be blind about the type of suture.

Placebo

Not used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Semnan University of Medical Sciences

Street address

Vice chancellor for research, Setad building, Basij Blvd

City

Semnan

Province

Semnan

Postal code

35198-99951

Approval date

2018-10-17, 1397/07/25

Ethics committee reference number

IR.SEMUMS.REC.1397.157

Health conditions studied

1

Description of health condition studied

Mandibular fully impacted third molar surgery

ICD-10 code

K01.1

ICD-10 code description

Impacted teeth

2

Description of health condition studied

Postoperative facial swelling after mandibular fully impacted third molar surgery

ICD-10 code

R22.9

ICD-10 code description

Localized swelling, mass and lump, unspecified

3

Description of health condition studied

Trismus after mandibular fully impacted third molar surgery

ICD-10 code

S03.4

ICD-10 code description

Sprain of jaw

4

Description of health condition studied

Dehiscence after mandibular fully impacted third molar surgery

ICD-10 code

T81.31

ICD-10 code description

Disruption of external operation (surgical) wound, not elsewhere classified

5

Description of health condition studied

Dry socket after mandibular fully impacted third molar surgery

ICD-10 code

K10.3

ICD-10 code description

Alveolitis of jaws

Primary outcomes

1

Description

pain

Timepoint

Before starting surgery, it is assessed that there is no pain at the surgical site. pain of the patient 12 hours after surgery and on days 1, 2, 3, 4, 5, 6 after surgery will be measured.

Method of measurement

Visual Analogue Scale

Secondary outcomes

1

Description

Swelling

Timepoint

Before starting surgery, 24 hours after surgery and the days 3rd and 7th after surgery

Method of measurement

Facial width will be measured by ticking three facial distances on a thread and then measuring it by a ruler. Horizontal: The distance between the corners of the lip to the attachment of the ear lobe on same side and the

thread is subject to the cheeks. Vertical: The distance between the outer eye cantos and the mandibular angle of the same side and the thread is subject to the cheeks. Oblique: The distance from the corners of the lip and the mandibular angle of the same side and the thread is subject to the possible facial swelling (post-operative swelling)

2

Description

Maximal oral opening rate

Timepoint

Before starting surgery, 24 hours after surgery and the days 3rd and 7th after surgery

Method of measurement

It will be calculated by measuring the distance between the central margin of the upper jaw and the central edge of the lower jaw (or left in the absence of the right central) by the digital caliper (Mitutoyo, Illinois, USA).

3

Description

Dry socket

Timepoint

24 hours after surgery and the days 3rd and 7th after surgery

Method of measurement

Blums criteria

4

Description

Dehiscence

Timepoint

24 hours after surgery and the days 3rd and 7th after surgery

Method of measurement

The self-opening of sutures without patient's intervention that define with observations

5

Description

Clinical attachment loss in 7th tooth distal

Timepoint

Before starting surgery, six months after surgery

Method of measurement

The distance between CEJ and the pocket bottom will be measured by william probe (Juya Instruments PVT, Tehran, Iran)

6

Description

The 7th tooth distal pocket depth

Timepoint

Before starting surgery, six months after surgery

Method of measurement

The distance between the gingival margin and pocket bottom will be measured by the william probe (Juya Instruments PVT, Tehran, Iran)

Intervention groups

1

Description

Intervention group: first the thread will be twisted around the needle holder in clockwise direction A) Then, with the aid of the needle holder the middle of the loose thread is taken and passed around the needle holder through the twisted thread to form a loop B) The needle holder is passed through the loop into the free end of the thread and pulled to complete the knot C) same process is repeated for the second time

Category

Treatment - Surgery

2

Description

Control group: A suture will be made on the releasing incision and two sutures will be implemented on 7th tooth distal and mesial papillae with a common knot.

Category

Treatment - Surgery

Recruitment centers

1

Recruitment center

Name of recruitment center

Semnan Dental School

Full name of responsible person

Mohammad Esmaeelnejad

Street address

Faculty of Dentistry, Semnan University of Medical Sciences, Hefdah Shahrivar St., Saadi Square

City

Semnan

Province

Semnan

Postal code

3519-99951

Phone

+98 23 3333 4657

Email

Esmaeelnejad@gmail.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Research center of Semnan Dental School

Full name of responsible person

Dr. Nazila Ameli

Street address

Research center, Semnan Dental School, 17 Shahrivar Blvd

City

Semnan

Province

Semnan

Postal code

35198-99951

Phone

+98 23 3333 4657

Email

Esmaeelnejad@gmail.com

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Research center of Semnan Dental School

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Semnan University of Medical Sciences

Full name of responsible person

Mohammad Esmaeelnejad

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Oral and maxillofacial surgery

Street address

Semnan Dental School, 17 Shahrivar Blvd

City

Semnan

Province

Semnan

Postal code

35198-99951

Phone

+98 23 3333 4657

Email

Esmaeelnejad@gmail.com

Person responsible for scientific inquiries

Contact

Name of organization / entity

Semnan University of Medical Sciences

Full name of responsible person

Mohammad Esmaeelnejad

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Dentistry

Street address

Semnan Dental School, 17 Shahrivar Blvd

City

Semnan

Province

Semnan

Postal code

35198-99951

Phone

+98 23 3333 4657

Email

Esmaeelnejad@gmail.com

Person responsible for updating data

Contact

Name of organization / entity

Semnan University of Medical Sciences

Full name of responsible person

MohammadHosein Mansourian

Position

Student

Latest degree

A Level or less

Other areas of specialty/work

Dentistry

Street address

No. 24,Shahid Ghavampour Ave.,North Sheikhbahaei Ave.,

City

Tehran

Province

Tehran

Postal code

1995763453

Phone

+98 21 8821 4780

Fax

Email

mh.mansourian95@gmail.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available