

Clinical Trial Protocol

Iranian Registry of Clinical Trials

17 Sep 2026

Comparison of the effect of continuous infusion of bupivacaine through intra-incisional catheter with control group for postoperative pain relieve in patients with intertrochanteric fracture

Protocol summary

Study aim

Comparison of the effect of continuous infusion of bupivacaine through intra-incisional catheter with control group for postoperative pain relieve in patients with intertrochanteric fracture

Design

Parallel Randomized Controlled Clinical Trial

Settings and conduct

At the end of the operation in the intervention group, an intra-incisional catheter will be placed under the fascial layer by surgeon, and after injection of 30 ml 0.25% bupivacaine it will be attached to an elastomeric pump containing 0.25% bupivacaine running 6 ml per hour. In control group no catheter will be placed. In both groups, an IV PCA morphine (0.5 mg/ml) pump will be run by the following setup: 2 ml bolus dose, 7 minutes lockout interval, no background infusion

Participants/Inclusion and exclusion criteria

Inclusion Criteria: Patients who scheduled to undergo hip nailing surgery and can tolerate surgery under spinal anesthesia, ASA I-II, 50-85 years of age. Exclusion Criteria: Uncontrolled seizure, Local infection, coagulopathy, thrombocytopenia, history of hepatic and renal functional impairment, narcotic addiction, allergy to drugs which are going to be used in this study, history of psychosomatic pain, psychiatric diseases, Diabetic neuropathy, 2nd and 3rd degree A-V block.

Intervention groups

In intervention group intra-incisional catheter will be placed for continuous bupivacaine infusion. Patients in both groups will be attached to IV morphine PCA pumps for better pain relieve.

Main outcome variables

Pain intensity during the first postoperative day; total morphine administered during the first postoperative day

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20100127003213N8**

Registration date: **2020-03-16, 1398/12/26**

Registration timing: **registered_while_recruiting**

Last update: **2020-03-16, 1398/12/26**

Update count: **0**

Registration date

2020-03-16, 1398/12/26

Registrant information

Name

Arash Farbood

Name of organization / entity

Shiraz University of Medical Sciences

Country

Iran (Islamic Republic of)

Phone

+98 71 1233 7636

Email address

farboda@sums.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2019-03-21, 1398/01/01

Expected recruitment end date

2020-04-20, 1399/02/01

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title
Comparison of the effect of continuous infusion of bupivacaine through intra-incisional catheter with control group for postoperative pain relieve in patients with intertrochanteric fracture

Public title
Effect of intra-incisional bupivacaine infusion in intertrochanteric fracture

Purpose
Prevention

Inclusion/Exclusion criteria
Inclusion criteria:
Patients who scheduled to undergo hip nailing surgery and can tolerate surgery under spinal anesthesia American Society of Anesthesiologists (ASA) I-II 50-85 years of age
Exclusion criteria:
Uncontrolled seizure Local infection Coagulopathy Thrombocytopenia History of hepatic and renal function impairment narcotic addiction Allergy to drugs which are going to be used in this study History of Psychosomatic pain Psychiatric diseases Diabetic neuropathy 2nd and 3rd degree A-V block

Age
From **50 years** old to **85 years** old

Gender
Both

Phase
2-3

Groups that have been masked
No information

Sample size
Target sample size: **48**

Randomization (investigator's opinion)
Randomized

Randomization description
Patients will be allocated into intervention and control groups by block randomization using randomization.com website. This will randomly generate a sequence of 12 blocks, 4 patients in each. The name of the patients' group will be concealed in separate envelopes which are identified by consecutive numbers related to the patient's entrance to the study.

Blinding (investigator's opinion)
Not blinded

Blinding description

Placebo
Not used

Assignment
Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Shiraz University of Medical Sciences

Street address

Vice chancellor of research, 7th floor of central building of Shiraz University of Medical Sciences, Zand street

City

Shiraz

Province

Fars

Postal code

7134844119

Approval date

2019-04-27, 1398/02/07

Ethics committee reference number

IR.SUMS.MED.REC.1398.099

Health conditions studied

1

Description of health condition studied

Femur fracture

ICD-10 code

S72.14

ICD-10 code description

Intertrochanteric fracture of femur

Primary outcomes

1

Description

Pain intensity

Timepoint

Every 15 minutes after surgery (15, 30, 45, 60 minutes) in recovery room, and every 2 hours after transfer to ward .

Method of measurement

Numerical Rating Scale of pain

2

Description

total morphine administration

Timepoint

total dose during the first postoperative 24 hours

Method of measurement

dose summation reported in milligrams

Secondary outcomes

1

Description

Patient's global satisfaction

Timepoint

Discharge from the ward

Method of measurement

A 5-point Likert scale: completely satisfied (= 5), partially satisfied (= 4), neutral (= 3), partially dissatisfied (= 2) and completely dissatisfied (= 1)

2

Description

Ease of ambulation

Timepoint

During admission

Method of measurement

Cumulated Ambulation Score

3

Description

Time to first request for analgesia

Timepoint

Time to first request, measured from termination of the procedure and recovery room entry

Method of measurement

minute

4

Description

Drug side effects (nausea & vomiting, respiratory depression, urinary retention, drowsiness, pruritus, hypotension, tinnitus)

Timepoint

Every four hours for 24 hours

Method of measurement

Occurrence frequency

Intervention groups

1

Description

Interventions group 1: At the end of the surgery an intra-incisional catheter (Infiltralong 600, PAJUNK, Geisingen, Germany) will be placed subfascially which will be connected to a 100 mL elastomeric pump, contained 0.25% bupivacaine by the rate of 6 ml per hour, after a 30 mL bolus dose of 0.25% bupivacaine (Bupivacaine® Mylan, 100 mg/20 ml vial, Delpharm, France). The intra-incisional catheter will be disconnected after 36 hours.

Category

Prevention

2

Description

Control group: The surgeon does not insert an intra-incisional catheter (Infiltralong manufacturing company. Pajunk of Germany) before the surgical wound is placed under the fascia layer. After the bolus injection of 2 ml of morphine sulfate (Aburaihan Pharmaceutical Co. Iran,) the pump settings are as follows: 20 ml of morphine sulfate at a concentration of 0.5 mg / ml in normal saline solution infused with infusion pump. Minimum injection interval: 7 minutes, maximal injection every 4 hours: 40

ml, and continued for 24 hours. infusing morphine, intravenously.

Category

Prevention

Recruitment centers

1

Recruitment center

Name of recruitment center

Chamran Hospital

Full name of responsible person

Sanaz Abbasi

Street address

Chamran Hospital, Chamran Boulevard

City

Shiraz

Province

Fars

Postal code

1564471948

Phone

+98 71 3624 0101

Email

chamranhosp@sums.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Shiraz University of Medical Sciences

Full name of responsible person

Dr Yones Ghasemi

Street address

Vice chancellor of research, 7th floor of central building of Shiraz University of Medical Sciences, Zand street

City

Shiraz

Province

Fars

Postal code

7134844119

Phone

+98 71 3647 4270

Email

sacrc@sums.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Shiraz University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
Academic

Person responsible for general inquiries

Contact

Name of organization / entity
Shiraz University of Medical Sciences
Full name of responsible person
Sanaz Abbasi
Position
Anesthesiology Resident
Latest degree
Medical doctor
Other areas of specialty/work
Anesthesiology
Street address
Anesthesiology Department, Faghihi Hospital, Zand
Street
City
Shiraz
Province
Fars
Postal code
7134844119
Phone
+98 71 3647 4270
Email
sanaz_med65@yahoo.com

Person responsible for scientific inquiries

Contact

Name of organization / entity
Shiraz University of Medical Sciences
Full name of responsible person
Arash Farbood
Position
Anesthesiologist/Pain Fellowship
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
Anesthesiology Department, Faghihi Hospital, Zand
Street
City
Shiraz

Province
Fars
Postal code
1564471948
Phone
+98 36474270
Email
arashfarbood@yahoo.com

Person responsible for updating data

Contact

Name of organization / entity
Shiraz University of Medical Sciences
Full name of responsible person
Arash Farbood
Position
Anesthesiologist/Pain Fellowship
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
Anesthesiology Department, Faghihi Hospital, Zand
Street
City
Shiraz
Province
Fars
Postal code
1564471948
Phone
+98 36474270
Email
arashfarbood@yahoo.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

No - There is not a plan to make this available

Justification/reason for indecision/not sharing IPD

It is against our policies.

Study Protocol

No - There is not a plan to make this available

Statistical Analysis Plan

No - There is not a plan to make this available

Informed Consent Form

No - There is not a plan to make this available

Clinical Study Report

No - There is not a plan to make this available

Analytic Code

No - There is not a plan to make this available

Data Dictionary

No - There is not a plan to make this available