

Clinical Trial Protocol

Iranian Registry of Clinical Trials

17 Jun 2026

Comparison effect of two approach for Improvement of patients Conditions in Recovery Using of oral or interavenous Glucose Pretreatment in Laparotomy surgeries

Protocol summary

Study aim

Comparison effect of oral or intravenous glucose pretreatment for improvement of recovery stay compared with control group

Design

In this double-blind randomized controlled clinical trial, faze 3 study with parallel groups, 90 patients undergoing laparotomy were randomly divided into 3 groups of 30 divided into the first group of intravenous glucose, the oral glucose guided group and the third group (control group) oral normal saline, 6 hours after fasting preoperatively were prescribed and recovery conditions were compared in three groups.

Settings and conduct

This study was carried out on patients undergoing laparotomy in Alzahra hospital in Isfahan in 1399. This study is done in two sucrose methods. Patients are unaware of the type of solution they are given and no explanation is given about the type of solution being given or injected. The solution is also prescribed by the patient's presenter, but the information is collected by an operating room staff who is not in the study.

Participants/Inclusion and exclusion criteria

Inclusion criteria:: Patients are candidates for laparotomy surgery, age range 18 to 65 years, physical health status 1 and 2. Non-entry criteria: Patients with diabetes and reflex, patients who need to take anti emetic drugs before surgery.

Intervention groups

In this study 90 patients undergoing laparotomy were divided into 3 groups of 30 each. All three groups received 10 ml/kg of serum 1/3-2/3 at the time of preoperative fasting, and 6 hours later, the first group received 1 g / kg glucose in 50 ml of normal saline intravenously injections, the second group 1 g / kg glucose in 50 ml of saline orally and the third group receive 50 ml of normal saline orally.

Main outcome variables

Postoperative pain, length of stay in recovery, nausea, vomiting

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20130311012782N55**

Registration date: **2021-07-10, 1400/04/19**

Registration timing: **retrospective**

Last update: **2021-07-10, 1400/04/19**

Update count: **0**

Registration date

2021-07-10, 1400/04/19

Registrant information

Name

Ali Mehrabi kushki

Name of organization / entity

Isfahan University of Medical Sciences

Country

Iran (Islamic Republic of)

Phone

+98 31 3629 1510

Email address

mehrabi@mui.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2020-08-22, 1399/06/01

Expected recruitment end date

2021-03-18, 1399/12/28

Actual recruitment start date

empty
Actual recruitment end date
empty
Trial completion date
empty

Scientific title
Comparison effect of two approach for Improvement of patients Conditions in Recovery Using of oral or interavenous Glucose Pretreatment in Laparotomy surgeries

Public title
Comparison effect of oral glucose administration and preoperative injection on patient status in recovery

Purpose
Prevention

Inclusion/Exclusion criteria

Inclusion criteria:

Patient undergoing laparotomy Patient's consent to participate in the study American Society of Anesthesiologists(ASA) : 1 or 2

Exclusion criteria:

Body Mass Index 30 or above diabetes reflux Requires preoperative Antiemetic medication

Age
From **18 years** old to **65 years** old

Gender
Both

Phase
3

Groups that have been masked

- Participant
- Outcome assessor

Sample size
Target sample size: **90**

Randomization (investigator's opinion)
Randomized

Randomization description
Randomization of patients is done using Random Allocation Aoft ware(RAS). The total sample size (90 people) is entered into the software and then the number of study groups (3 groups) is determined. The software output contains a list that distributes numbers 1 to 90 randomly in three columns. Patients are divided into groups according to the time list of admission to the hospital to reach the required sample size in each group.

Blinding (investigator's opinion)
Double blinded

Blinding description
This study is a double blind study. Patients are unaware of the type of solution they receive. The solution used by the researcher is also prescribed for patients, but the information is collected by one of the operating room personnel who is unaware of the type of solution being received.

Placebo
Used

Assignment
Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Isfahan University of Medical Sciences

Street address

Research faculty, Isfahan University of Medical Sciences, Hezarjerib street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Approval date

2019-02-09, 1397/11/20

Ethics committee reference number

IR.MUI.MED.REC.1398.010

Health conditions studied

1

Description of health condition studied

Recovery condition

ICD-10 code

K91.86

ICD-10 code description

cholecystectomy

Primary outcomes

1

Description

post operative pain

Timepoint

during recovery stay

Method of measurement

Visual Analog Scale

2

Description

stay in recovery time

Timepoint

End of recovery

Method of measurement

Calculates the time interval of entry to exit recovery in minutes

3

Description

nausea

Timepoint

During recovery stay

Method of measurement

Ask the patient

4**Description**

Vomitting

Timepoint

During recovery stay

Method of measurement

Ask the patient

Secondary outcomes

empty

Intervention groups**1****Description**

Intervention group 1: 10 ml/ kg of serum 1/3-2/3 at first fasting and after 6 hours, 1 g / kg glucose dissolved in 50 ml of normal saline is injected intravenously.

Category

Prevention

2**Description**

Intervention group 2: 10 ml/ kg of serum 1/3-2/3 at first fasting and after 6 hours, 1 g / kg glucose dissolved in 50 ml of normal saline is orally prescribed.

Category

Prevention

3**Description**

Control group: 10 ml/ kg of serum 1/3-2/3 at first fasting and after 6 hours, 50 ml of normal saline is orally prescribed.

Category

Prevention

Recruitment centers**1****Recruitment center****Name of recruitment center**

Alzahra hospital

Full name of responsible person

Ladan Salehfard

Street address

Department of Anesthesiology, Alzahra hospital,
Soffeh street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Phone

+98 31 3668 5555

Email

ladan.salehfard@gmail.com

Sponsors / Funding sources**1****Sponsor****Name of organization / entity**

Esfahan University of Medical Sciences

Full name of responsible person

Shaghayegh Haghjoo

Street address

Research faculty, Isfahan University of Medical
Sciences, Hezarjerib street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Phone

+98 31 3792 3070

Email

sh_haghjoo@med.mui.ac.ir

Grant name**Grant code / Reference number****Is the source of funding the same sponsor organization/entity?**

Yes

Title of funding source

Esfahan University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin**Type of organization providing the funding**

Academic

Person responsible for general inquiries**Contact****Name of organization / entity**

Esfahan University of Medical Sciences

Full name of responsible person

Taghi Hashemi

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Department of anesthesiology, Alzahra hospital,
Soffeh street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Phone

+98 31 3668 5555

Email

st_hashemi@med.mui.ac.ir

Person responsible for scientific inquiries

Contact

Name of organization / entity

Esfahan University of Medical Sciences

Full name of responsible person

Mitra Jabalameli

Position

Associate professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Department of Anesthesiology, Alzahra hospital,
Soffeh street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Phone

+98 31 3668 5555

Email

Jabalameli@med.mui.ac.ir

Person responsible for updating data

Contact

Name of organization / entity

Esfahan University of Medical Sciences

Full name of responsible person

Ali Mehrabi

Position

Statistical Consultant

Latest degree

Master

Other areas of specialty/work

Epidemiology

Street address

Research faculty, school of medicine, Isfahan
University of Medical Sciences, Hezarjerib street

City

Isfahan

Province

Isfahan

Postal code

8434193474

Phone

0098 31 3728081

Email

al.mehrabi@gmail.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

No - There is not a plan to make this available

Justification/reason for indecision/not sharing IPD

The project is owned by the state agency and cannot be published.

Study Protocol

No - There is not a plan to make this available

Statistical Analysis Plan

No - There is not a plan to make this available

Informed Consent Form

No - There is not a plan to make this available

Clinical Study Report

No - There is not a plan to make this available

Analytic Code

No - There is not a plan to make this available

Data Dictionary

No - There is not a plan to make this available