

Clinical Trial Protocol

Iranian Registry of Clinical Trials

01 Jul 2026

Comparison of luteal phase stimulation with follicular phase stimulation in poor ovarian response

Protocol summary

Study aim

Comparison of luteal phase stimulation with follicular phase stimulation in poor ovarian response

Design

Clinical trial with control group, with parallel groups, single-blind, randomized, phase 3 on 78 patients. Randomization will be done using 4 blocks designed with sealedenvelop.com software.

Settings and conduct

This study will be performed at Avicenna Infertility Center. Women are randomly divided into two groups. In the intervention group ovarian stimulation begins in the luteal phase and in the control group ovarian stimulation begins in the follicular phase. In both groups, GnRH antagonist is administered when the largest follicle is > 12 mm. Stimulation of ovulation using a GnRH agonist with 2 HCGs when at least one follicle > 18 mm is seen will be prescribed. Oocyte collection will be done by transvaginal ultrasound 36 hours later. Fertilization will take place after oocyte retrieval. The embryos are then frozen on the third or fifth day, and in another cycle, the uterus is prepared for embryo transfer and the embryos are transferred. Pregnancy test is done 14 days after embryo transfer. Chemical and clinical pregnancy rates will be assessed.

Participants/Inclusion and exclusion criteria

- History of an ovulation stimulation cycle with less than 4 oocytes - AMH < 1.1 ng/ml or AFC < 5 - Total number of motile sperm at least 15 million - Normal uterus in ultrasound hysterosalpingography - Regular periods of 21-35 days - Indication for IVF - Compliance to participate and collaborate with study protocol - Signing informed consent

Intervention groups

Intervention group: ovarian stimulation in the luteal phase will be started on days 15-16 of the cycle. Stimulation will be performed using HCG. Control group: ovarian stimulation in the follicular phase is initiated at the beginning of the cycle. Stimulation will be performed

using HCG.

Main outcome variables

Number of metaphase II oocytes

General information

Reason for update

The update is to change the sample size.

Acronym

IRCT registration information

IRCT registration number: **IRCT20210405050852N1**

Registration date: **2021-06-10, 1400/03/20**

Registration timing: **prospective**

Last update: **2023-03-01, 1401/12/10**

Update count: **1**

Registration date

2021-06-10, 1400/03/20

Registrant information

Name

Mozhgan Vahabi Dastjerdi

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 2264 4701

Email address

vahabi.mozhgan@yahoo.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2021-07-01, 1400/04/10

Expected recruitment end date

2021-12-31, 1400/10/10

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparison of luteal phase stimulation with follicular phase stimulation in poor ovarian response

Public title

Effect of luteal phase stimulation with follicular phase stimulation in poor ovarian response

Purpose

Treatment

Inclusion/Exclusion criteria**Inclusion criteria:**

History of an ovulation stimulation cycle with less than 4 oocytes AMH<1.1 ng/ml or AFC<5 The total number of motile sperm is at least 15 million Normal uterus in ultrasound and hysterosalpingography Regular periods of 21-35 days Indication for IVF Ability to participate and collaborate with study protocol Signature informed consent Under 40 years

Exclusion criteria:

History of ovarian surgery History of pelvic cytotoxic radiation exposure due to malignancy History of recurrent miscarriage Grade III or IV endometriosis Concomitant uterine pathology (adenomyosis, submucosal myoma, Asherman syndrome) Autoimmune disease

Age

From **18 years** old to **39 years** old

Gender

Female

Phase

3

Groups that have been masked

- Participant

Sample size

Target sample size: **78**

Randomization (investigator's opinion)

Randomized

Randomization description

The Block randomization method was designed by an epidemiologist using STATA software version 13 and the number of blocks considered is 4. The random allocation list for patients was solely available to the epidemiologist. In order to hide the random allocation process, a total of 78 envelopes were prepared, and only the methodologist has been aware of the table of random numbers. When the doctor declared the patient's eligibility, the methodologist provided the doctor with the envelope. The group was selected based on the type of group mentioned in the envelope.

Blinding (investigator's opinion)

Single blinded

Blinding description

This is a single-blind study. The participants don't aware about grouping name and type of treatment in each group. In order to blind the patients participating in this study, all conditions will be the same between the two

groups, so patients in the intervention and control groups will refer to the center on ovulation stimulation days in both groups.

Placebo

Not used

Assignment

Parallel

Other design features**Secondary Ids**

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Ethics Committee of Avicenna Research Institute (ARI)

Street address

No. 97, at the corner of the Yakhchal st, Shariati st, Avicenna Infertility Clinic

City

Tehran

Province

Tehran

Postal code

1941913114

Approval date

2021-03-09, 1399/12/19

Ethics committee reference number

IR.ACECR.AVICENNA.REC.1400.002

Health conditions studied**1****Description of health condition studied**

Poor ovarian response

ICD-10 code

E28

ICD-10 code description

Ovarian dysfunction

Primary outcomes**1****Description**

Number of metaphase II oocyte

Timepoint

End of ovarian stimulation

Method of measurement

Observation of metaphase 2 oocytes extracted by the embryologist

Secondary outcomes

1

Description

Chemical pregnancy rate

Timepoint

14 days after transfer

Method of measurement

Beta-Human Chorionic Gonadotropin

2

Description

Clinical pregnancy

Timepoint

Pregnancy 7th weeks

Method of measurement

Detect of fetal heart beat on ultrasound

3

Description

Total number of embryos

Timepoint

After in vitro fertilization

Method of measurement

18-20 hours after in vitro fertilization

Intervention groups

1

Description

In this study, to compare the effect of luteal phase stimulation with follicular phase stimulation in patients with poor ovarian response, in the luteal phase group on days 15 to 16 of the menstrual cycle and when spontaneous ovulation is confirmed by transvaginal ultrasound and patients have one follicle at least 8 mm, start with 300 units of FSH plus 150 to 225 units of HMG plus 10 mg of medroxyprogesterone daily. Medroxyprogesterone is used to delay menstruation and stop the release of oocytes during menstruation. GnRH antagonist is given when the largest follicle is > 12 mm. Sono examination will be performed every 24 to 72 hours. Blood hormones will also be tested by checking estradiol and progesterone levels on the day of ovulation stimulation. Ovulation stimulation will be administered using a GnRH agonist with 2 Human chorionic gonadotropins when at least one follicle <18 mm is seen. Oocyte collection will be done by transvaginal ultrasound 36 hours later. Fertilization will take place after oocyte retrieval. Successful fertilization is diagnosed when two pronuclei are seen about 18-20 hours after insemination. The embryos are then frozen on the third or fifth day, and in another cycle, the uterus is prepared for embryo transfer and the embryos are transferred. 14 days after the transfer of the fetus, the pregnancy test will be requested and the rate of chemical and clinical pregnancy will be evaluated.

Category

Treatment - Drugs

2

Description

Control group: In patients in the control group (follicular phase stimulation group), ovarian stimulation begins on the second or third day of the cycle with 300 units of FSH plus 150 to 225 units of HMG. GnRH antagonist is given when the largest follicle is > 12 mm. Ultrasound examination will be performed every 24 to 72 hours. Blood hormone analysis will also be done by checking estradiol and progesterone levels on the day of ovulation stimulation, and ovulation stimulation by GnRH agonist along with 2 Human chorionic gonadotropins will be administered when at least one follicle <18 mm is seen. Fertilization will take place after oocyte harvest. Successful fertilization is diagnosed when two pronuclei are seen about 18-20 hours after inoculation. The embryos are then frozen on the third or fifth day, and in another cycle, the uterus is prepared for embryo transfer and the embryos are transferred. 14 days after the transfer of the fetus, the pregnancy test will be requested and the rate of chemical and clinical pregnancy will be evaluated.

Category

Treatment - Drugs

Recruitment centers

1

Recruitment center

Name of recruitment center

Avicenna infertility clinic

Full name of responsible person

Simin Zafardoost

Street address

No. 97, at the corner of the Yakhchal st, Shariati st, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1941913118

Phone

+98 21 2264 4706

Fax

+98 21 2264 4754

Email

siminzafardoost@yahoo.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Avicenna University Jihad Infertility Center Avicenna

Full name of responsible person

Ramin Ghahremanzadeh

Street address

No. 97, at the corner of the Yakhchal st, Shariati st,

Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1941913114

Phone

+98 21 2264 4706

Fax

+98 21 2264 4754

Email

Vahabi.mozhgan@yahoo.com

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Avicenna University Jihad Infertility Center Avicenna

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Avicenna infertility clinic

Full name of responsible person

Mozhgan Vahabi Dastjerdi

Position

Infertility Fellowship

Latest degree

Specialist

Other areas of specialty/work

Gynecology and Obstetrics

Street address

No. 97, at the corner of the Yakhchal st, Shariati st, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1941913114

Phone

+98 21 2264 4706

Fax

+98 21 2264 4754

Email

Vahabi.mozhgan@yahoo.com

Person responsible for scientific inquiries

Contact

Name of organization / entity

Avicenna infertility center

Full name of responsible person

Mozhgan Vahabi Dastjerdi

Position

Infertility Fellowship

Latest degree

Specialist

Other areas of specialty/work

Gynecology and Obstetrics

Street address

No. 97, at the corner of the Yakhchal st, Shariati st, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1941913114

Phone

+98 21 2264 4706

Fax

+98 21 2264 4754

Email

vahabi.mozhgan@yahoo.com

Person responsible for updating data

Contact

Name of organization / entity

Avicenna Infertility center

Full name of responsible person

Mozhgan Vahabi Datjerdi

Position

Infertility Fellowship

Latest degree

Specialist

Other areas of specialty/work

Gynecology and Obstetrics

Street address

No. 97, at the corner of the Yakhchal st, Shariati st, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1941913114

Phone

+98 21 2264 4706

Fax

+98 21 2264 4754

Email

vahabi.mozhgan@yahoo.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Yes - There is a plan to make this available
Study Protocol
Yes - There is a plan to make this available
Statistical Analysis Plan
Yes - There is a plan to make this available
Informed Consent Form
Yes - There is a plan to make this available
Clinical Study Report
Yes - There is a plan to make this available
Analytic Code
Yes - There is a plan to make this available
Data Dictionary
Undecided - It is not yet known if there will be a plan to make this available
Title and more details about the data/document
Comparison of luteal phase stimulation with follicular

phase stimulation in poor ovarian response
When the data will become available and for how long
Up to one year after the publication of the article
To whom data/document is available
Other researchers and physicians
Under which criteria data/document could be used
Use in meta-analysis studies
From where data/document is obtainable
Avicenna infertility clinic, Dr. Mozghan Vahabi datjerdi,
00982122644706
What processes are involved for a request to access data/document
Email to the responsible author, Dr. Mozghan vahabi
Dastjerdi
Comments