

Clinical Trial Protocol

Iranian Registry of Clinical Trials

09 Jun 2026

Assessment of Oxycodone and Mebeverin mixed-effects on colonoscopy quality in patients referring to Urmia Imam Khomeini hospital

Protocol summary

Study aim

Determining the combined effect of oxycodone and Mebeverine on colonoscopy quality in patients referred to Imam Khomeini Hospital

Design

Clinical trial with control group, with parallel groups, double-blind, randomized by random number table, phase three on 125 patients.

Settings and conduct

125 people referring to the colonoscopy ward of Imam Khomeini Educational and Medical Center in Urmia, if they have inclusion criteria and do not have exclusion criteria. after the randomization by the table of lucky numbers and blinding so that if the number is out between 1 to 62 The patient will be in the intervention group and will receive the drug (oxycodone and Mebeverine) and if the number is between 63 to 125 will be in the control group and will receive the placebo (vitamin B6 tablets and starch capsules) one hour before colonoscopy.

Participants/Inclusion and exclusion criteria

Anyone over the age of 18 who is suspected of having intestinal polyps, unexplained gastrointestinal bleeding, and intestinal disease will undergo a colonoscopy with the inclusion criteria of do not take antispasmodics drugs, complete mental health for informed consent, insensitivity to oxycodone and Mebeverine and no history of abdominal surgery. Non-entry conditions: History of allergy to any type of drug and irritable bowel syndrome.

Intervention groups

In the intervention group, one oxycodone tablet and two Mebeverine capsules will be given to the patient one hour before colonoscopy by someone other than the endoscopist. In the control group, a vitamin B6 tablet will be given to the patient one hour before colonoscopy along with two starch-containing capsules.

Main outcome variables

Total procedure time, pain, time to reach the cecum

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20201204049599N3**

Registration date: **2021-04-22, 1400/02/02**

Registration timing: **prospective**

Last update: **2021-04-22, 1400/02/02**

Update count: **0**

Registration date

2021-04-22, 1400/02/02

Registrant information

Name

Mohammad Reza Pashaee

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 44 3198 8001

Email address

pashaee.m@umsu.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2021-04-30, 1400/02/10

Expected recruitment end date

2021-05-22, 1400/03/01

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Assessment of Oxycodone and Mebeverin mixed-effects on colonoscopy quality in patients referring to Urmia Imam Khomeini hospital

Public title

Assessment of Oxycodone and Mebeverin mixed-effects on colonoscopy quality

Purpose

Treatment

Inclusion/Exclusion criteria

Inclusion criteria:

Age over 18 years Do not take antispasmodics Suspicion of intestinal polyps bowel diseases Unknown gastrointestinal bleeding

Exclusion criteria:

Irritable Bowel Syndrome Previous abdominal surgeries Lack of mental health and full consciousness Allergy to oxycodone Allergy to Mebeverine Taking anti-anxiety and painkillers History of allergy to any drug

Age

From **18 years** old

Gender

Both

Phase

3

Groups that have been masked

- Participant
- Investigator

Sample size

Target sample size: **125**

Randomization (investigator's opinion)

Randomized

Randomization description

Randomization method is simple randomization. the researcher will divide patients by using table of random numbers. First, each patient will be assigned a three-digit code according to the date and time of the clinic visit. The first patient will have code 001 and the last patient will have code 125. Then in the table of random numbers without looking, a location will be randomly selected. The first two digits will be examined. If the number is less than or equal to 125, it will be selected, otherwise we will continue moving in the direction of the line and other numbers will be checked. If it is a duplicate number, it will be ignored and another number from the table will be checked. We continue this process until 62 codes are selected. The first 62 codes will be for patients in the intervention group (patients who will receive drugs) and the remaining 63 codes will be for the control group (patients who will receive placebo). Thus, patients are randomly divided into two groups.

Blinding (investigator's opinion)

Double blinded

Blinding description

The study will be two-way blind. The patient and the endoscopist will be unaware that the patient belongs to the case or control group. Pills and capsules in the same shape, size and color will be given to patients in both groups. In the intervention group, one oxycodone tablet, which is small and white, and two Mebeverine capsules, which are white, will be given to the patient one hour

before colonoscopy by someone other than the endoscopist. In the control group, a vitamin B6 tablet, which is similar in shape, color and size to oxycodone and is completely safe, will be given to the patient one hour before colonoscopy, along with two starch-containing capsules that resemble Mebeverine capsules.

Placebo

Used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Urmia University of Medical Sciences

Street address

Urmia University of Medical Sciences; Resalat boulevard

City

Urmia

Province

West Azarbaijan

Postal code

5714783734

Approval date

2020-09-30, 1399/07/09

Ethics committee reference number

IR.UMSU.REC.1399.261

Health conditions studied

1

Description of health condition studied

colonoscopy

ICD-10 code

K92.2

ICD-10 code description

Gastrointestinal hemorrhage, unspecified

Primary outcomes

1

Description

Total procedure time

Timepoint

During endoscopy

Method of measurement

minute, clock

2

Description

pain

Timepoint

During endoscopy

Method of measurement

Score based on visual analog scale

3

Description

Time to reach Cecum

Timepoint

During endoscopy

Method of measurement

minute , clock

Secondary outcomes

empty

Intervention groups

1

Description

Intervention group: One hour before colonoscopy, patients in this group will be given 15 mg of oral Oxycodone tablet and 270 mg of oral Mebeverine capsule .

Category

Treatment - Drugs

2

Description

Control group : In the control group, a vitamin B6 tablet, which is similar in shape, color and size to oxycodone and is completely safe, will be given to the patient one hour before colonoscopy, along with two starch-containing capsules that resemble Mebeverine capsules.

Category

Placebo

Recruitment centers

1

Recruitment center

Name of recruitment center

Imam Khomeini Hospital

Full name of responsible person

Mohammad Reza Pashae

Street address

Imam Khomeini Hospital; Ershad boulevard

City

Urmia

Province

West Azarbaijan

Postal code

5714783734

Phone

+98 44 3398 1002

Email

pashae.m@umsu.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Oroumia University of Medical Sciences

Full name of responsible person

Dr. Iraj Mohebbi

Street address

Urmia University of Medical Sciences; Resalat boulevard

City

Urmia

Province

West Azarbaijan

Postal code

5714783734

Phone

+98 44 3223 4897

Email

i.mohebbi@umsu.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Oroumia University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Oroumia University of Medical Sciences

Full name of responsible person

Mohammad Reza Pashae

Position

Associate professor

Latest degree

Subspecialist

Other areas of specialty/work

Internal Medicine

Street address

Imam Khomeini Hospital; Ershad boulevard

City

Urmia

Province
West Azarbaijan
Postal code
5714783734
Phone
+98 44 3198 8001
Fax
Email
pashae.m@umsu.ac.ir

Person responsible for scientific inquiries

Contact

Name of organization / entity
Oroumia University of Medical Sciences
Full name of responsible person
Mohammad Reza Pashae
Position
Associate professor
Latest degree
Subspecialist
Other areas of specialty/work
Internal Medicine
Street address
Imam Khomeini Hospital; Ershad boulevard
City
Urmia
Province
West Azarbaijan
Postal code
5714783734
Phone
+98 44 3198 8001
Fax
Email
pashae.m@umsu.ac.ir

Person responsible for updating data

Contact

Name of organization / entity
Oroumia University of Medical Sciences
Full name of responsible person

Mohammad Reza Pashae
Position
Associate professor
Latest degree
Subspecialist
Other areas of specialty/work
Internal Medicine
Street address
Imam Khomeini Hospital; Ershad boulevard
City
Urmia
Province
West Azarbaijan
Postal code
5714783734
Phone
+98 44 3198 8001
Fax
Email
pashae.m@umsu.ac.ir

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available