

Clinical Trial Protocol

Iranian Registry of Clinical Trials

30 Jun 2026

Evaluation of the effects of endometrial scratching on angiogenic and anti-angiogenic growth factors expression in unexplained repeated implantation failure (uRIF) patients

Protocol summary

Study aim

Determining the effect of endometrial scratching on the expression of angiogenic and anti-angiogenic growth factors

Design

A randomized clinical trial with control group, double blind, and two arm parallel group design of 20 patients. Randomization is performed using a computer-generated random assignment schedule for each patient. Sealed and numbered envelopes are used to conceal the treatment allocation until randomization.

Settings and conduct

Using a pipelle, endometrial scratching is performed by a gynecologist only in the intervention group in the clinic. In this study, the participating patients (creating the same conditions) and the researcher (through sample coding) will be blinded.

Participants/Inclusion and exclusion criteria

Inclusion criteria: Women with Unexplained repeated implantation failure(uRIF), 20 to 40 years, Body max index less (BMI) than 25, good response to previous ovulation stimulation, at least two embryos with good grade in the current cycle, normal uterus. Exclusion criteria: Endometrial thickness is less than 7 mm, congenital anomalies,myoma,endometrioma, adhesions, hydrosalpinx, previous uterine or ovarian surgery, severe male factor infertility, number of embryos less than 2 in the current cycle, diabetes, thyroid disease, any endocrine, genetic, infection or autoimmune disorder

Intervention groups

Intervention: In the intervention group, endometrial sampling is obtained twice by Pipelle [one in the follicular phase and the last in the luteal phase of the same cycle preceding the embryo transfer cycle. Control group: In the control group endometrial sampling will be done only in the luteal phase of the cycle preceding the embryo transfer cycle.

Main outcome variables

Determination of angiogenic and anti-angiogenic growth factor gene expression in endometrial specimens in the intervention group compared with the control group

General information

Reason for update

According to the previous report, the actual recruitment end date of hospitalization is the same as 01/26/1401, and the only date of trial completion date is according to the examination of clinical pregnancy (5 weeks after embryo transfer) and live birth (9 months of pregnancy)in the participant's plan has increased. In addition, according to the evaluation of the secondary variables of implantation rate and clinical pregnancy, the variables related to them that are evaluated are added. A little rewriting has been done in the way the inclusion and exclusion criteria are expressed.

Acronym

IRCT registration information

IRCT registration number: **IRCT20210316050723N1**
Registration date: **2021-07-12, 1400/04/21**
Registration timing: **registered_while_recruiting**

Last update: **2024-01-27, 1402/11/07**

Update count: **4**

Registration date

2021-07-12, 1400/04/21

Registrant information

Name

Samaneh Aghajanzpour

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 2356 2727

Email address

Recruitment status
Recruitment complete
Funding source

Expected recruitment start date

2021-06-20, 1400/03/30

Expected recruitment end date

2023-06-20, 1402/03/30

Actual recruitment start date

2021-06-20, 1400/03/30

Actual recruitment end date

2022-04-15, 1401/01/26

Trial completion date

2023-01-21, 1401/11/01

Scientific title

Evaluation of the effects of endometrial scratching on angiogenic and anti-angiogenic growth factors expression in unexplained repeated implantation failure (uRIF) patients

Public title

Evaluation of the effects of endometrial scratching on angiogenic and anti-angiogenic growth factors

Purpose

Other

Inclusion/Exclusion criteria

Inclusion criteria:

Infertile women with unexplained repeated implantation failure (unknown definite causes of RIF) failed to conceive after three or more embryo transfer cycles using high-quality transferred embryos (at least one blastocyst ET cycle) Age of 20 to 40 years Body mass index (BMI) less than 25 kg/m² Good response to stimulation of previous ovulation Having at least two embryos with good grade normal uterus results of hysterosalpingography (HSG) or hysteroscopy Participant consent was required to join in the study and to complete the consent form

Exclusion criteria:

The thickness of the endometrium on the day of Human chorionic gonadotropin hormone injection is less than 7 mm Women with congenital anomalies, intramural and subserosal myomas(larger than 5 cm) Submucosal myoma Women with endometrioma larger than or equal to 3 cm With adhesions With hydrosalpinx Have previous uterine or ovarian surgery Severe male factor infertility (sperm extraction from the testis, sperm freezing, sperm DNA fragmentation index equal to or above 16%) Patients with any specific medication If the number of available embryos is less than 2 in the current cycle Women with endometrial tuberculosis and those undergoing tuberculosis treatment Have a history of diabetes, thyroid disease, any endocrine, genetic, infection or autoimmune disorder Abnormal Pre-implantation Genetic Test (PGT) Results Any specific medication Failure to return the patient to prepare an endometrial sample Women with severe pain during obtaining of tissue samples or the possibility of infection

Age

From **20 years** old to **40 years** old

Gender

Female

Phase

2

Groups that have been masked

- Participant
- Investigator

Sample size

Target sample size: **20**

Actual sample size reached: **20**

Randomization (investigator's opinion)

Randomized

Randomization description

Consecutive sampling until the required sample size is reached. Women by Balanced Block Randomization method are randomly divided into 2 equal groups of 10 people (intervention group and control group). Block randomization method is designed by epidemiologist using STATA software version 13 and the number of blocks considered is 4. Envelopes are prepared for 20 people and inside each envelope is written the group in which the patient should be placed. The envelopes are prepared in a way that the writing inside is not clear. A nurse before the patient enters the operating room, removes the envelope and sends the patients in one of the two groups.

Blinding (investigator's opinion)

Double blinded

Blinding description

In this study, a doctor scratches the endometrium in the intervention group on days 8-11. The biopsy specimen is prepared in the luteal phase of the same cycle during days 19-23 by the same doctor or other. Therefore, in this study, it is not possible to blind the doctor. In order to blind the patients participating in this study, all the conditions will be the same between the two groups, so the patients of the control group also referred to the center on the day of the scratching (8-11) and due to the blinding of the study, all the sampling steps were done except endometrial scratching will be done for the control group as well as the intervention group. Then, biopsy samples are prepared from both groups in the luteal phase on days 19-23. The endometrial biopsy sample of both groups is sent to the laboratory, which does not know whether the tissue sample received is for the intervention group or the control group, and only checks it based on the received code (blinding of the researcher).

Placebo

Not used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Research Ethics Committees of school of medicine

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran
University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۳۹۶۱۴۵۳۵

Approval date

2021-06-12, 1400/03/22

Ethics committee reference number

IR.IUMS.FMD.REC.1400.147

Health conditions studied

1

Description of health condition studied

Unexplained repeated implantation failure (uRIF)

ICD-10 code

N97.9

ICD-10 code description

Female infertility, unspecified

Primary outcomes

1

Description

Evaluation of gene expression

Timepoint

In the endometrial sample obtained in the luteal phase during days 19-23

Method of measurement

Using PCR Array method and based on the copy number

Secondary outcomes

1

Description

Clinical pregnancy

Timepoint

5 weeks after embryo transfer

Method of measurement

Vaginal ultrasonography (observing the gestational sac and fetal heart rate)

2

Description

Implantation

Timepoint

5 weeks after embryo transfer

Method of measurement

The number of observed sacs divided by the number of

transferred embryos

3

Description

Total dose of gonadotropin (IU)

Timepoint

From ovarian stimulation to human chorionic gonadotropin (hCG) injection

Method of measurement

The number of international units (IU) received during the ovarian stimulation cycle

4

Description

Duration of ovarian stimulation (day)

Timepoint

From the start of gonadotropin administration to human chorionic gonadotropin (hCG) injection

Method of measurement

According to the patient's monitoring ultrasound chart (number of days receiving the medicine)

5

Description

Number of retrieved oocytes

Timepoint

On the day of oocyte pick-up

Method of measurement

The number of retrieved oocytes reported in the embryology chart

6

Description

Metaphase II oocytes

Timepoint

On the day of oocyte pick-up

Method of measurement

Based on the presence of the polar body reported by the embryologist

7

Description

Number of embryos

Timepoint

Three or five days after oocyte pick-up

Method of measurement

The number of cleavage embryos reported by the embryologist

8

Description

Embryo transfer cancelation

Timepoint

Three or five days after oocyte pick-up

Method of measurement

Participants with no embryo transfer

9

Description

Chemical pregnancy rate/Embryo transfer

Timepoint

2 weeks after embryo transfer

Method of measurement

Beta human chorionic gonadotropin (beta hCG) titer in blood serum by ELISA method

10

Description

Blighted ovum/Embryo transfer

Timepoint

5 weeks after embryo transfer

Method of measurement

Absence of fetus in gestational sac according to ultrasound report

11

Description

Ectopic pregnancy/Embryo transfer

Timepoint

5 weeks after embryo transfer

Method of measurement

According to the ultrasound report, there is a gestational sac outside the uterus with a positive Beta hCG titer

12

Description

Miscarriage rate/Embryo transfer

Timepoint

Pregnancy loss before 20 weeks

Method of measurement

Excretion of pregnancy remnants with vaginal bleeding (absence of heartbeat according to ultrasound report)

13

Description

Multiple Pregnancy (Twin)

Timepoint

5 weeks after embryo transfer

Method of measurement

Ultrasound report based on the number of gestational sacs with embryo

14

Description

Live birth rate/Embryo transfer

Timepoint

Time of delivery

Method of measurement

Birth of a live baby

15

Description

Fertilization rate

Timepoint

The day after the sperm injection

Method of measurement

number of oocytes with two pronuclei (2PN) divided by the number of injected oocytes

Intervention groups

1

Description

Intervention group: In the intervention group, endometrial sampling is obtained twice by Pipelle [one in the follicular phase (during 8-11 days and the last in the luteal phase (during 19-23 days) preceding the embryo transfer cycle. The endometrial scratching is induced with pipelle.

Category

Treatment - Other

2

Description

Control group: In the control group endometrial sampling will be done only in the luteal phase of the cycle preceding the embryo transfer cycle.

Category

Treatment - Other

Recruitment centers

1

Recruitment center

Name of recruitment center

Laleh Hospital

Full name of responsible person

Mehrdad Bakhtiyari

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran
University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۴۹۶۱۴۵۳۵

Phone

+98 21 8670 4567

Email

Mehr_bakhtiyari@yahoo.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Iran University of Medical Sciences

Full name of responsible person

Mehrdad Bakhtiyari

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran

University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۴۹۶۱۴۵۳۵

Phone

+98 21 8670 4567

Email

Mehr_bakhtiyari@yahoo.com

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Iran University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Iran University of Medical Sciences

Full name of responsible person

Mehrdad Bakhtiyari

Position

Assistant professor

Latest degree

Ph.D.

Other areas of specialty/work

Anatomy

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran
University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۴۹۶۱۴۵۳۵

Phone

+98 21 8670 4567

Email

mehr_bakhtiyari@yahoo.com

Person responsible for scientific inquiries

Contact

Name of organization / entity

Iran University of Medical Sciences

Full name of responsible person

Mehrdad Bakhtiyari

Position

Assistant professor

Latest degree

Ph.D.

Other areas of specialty/work

Anatomy

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran
University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۴۹۶۱۴۵۳۵

Phone

+98 21 8670 4567

Email

Mehr_bakhtiyari@yahoo.com

Person responsible for updating data

Contact

Name of organization / entity

Iran University of Medical Sciences

Full name of responsible person

Mehrdad Bakhtiyari

Position

Assistant professor

Latest degree

Ph.D.

Other areas of specialty/work

Anatomy

Street address

Tehran, Hemmat Highway next to Milad Tower, Iran
University of Medical Sciences

City

Tehran

Province

Tehran

Postal code

۱۴۴۹۶۱۴۵۳۵

Phone

+98 21 8670 4567

Email

Mehr_bakhtiyari@yahoo.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Yes - There is a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to make this available

Clinical Study Report

Yes - There is a plan to make this available

Analytic Code

Not applicable

Data Dictionary

Not applicable

Title and more details about the data/document

Clinical study report (published article)

When the data will become available and for how long

After the publication of the article

To whom data/document is available

Available to the public

Under which criteria data/document could be used

Scientific use by citing the source

From where data/document is obtainable

Dr. Mehrdad Bakhtiyari

What processes are involved for a request to access data/document

Request via e-mail

Comments