

Clinical Trial Protocol

Iranian Registry of Clinical Trials

18 Sep 2026

Comparison of efficacy of ultrasound guided pectoral nerve block (PECS II) versus serratus anterior block (SAB) on postoperative pain in breast cancer surgery

Protocol summary

Study aim

Comparison of efficacy of ultrasound guided pectoral nerve block (PECS II) versus serratus anterior block on postoperative pain in breast cancer surgery

Design

Clinical trial with control group, with parallel groups, One-blind, randomized, phase 3 on 300 patients. In order to randomize, the block randomization method will be used.

Settings and conduct

In this study, all patients who are candidates for breast cancer surgery at Rasoul Hospital in Tehran will be enrolled. Patients will be randomly divided into 2 groups based on size 4 blocks. The sample size for each study group is 40 people. A total of 80 patients will be examined. Patients and information analysts will be blind. The amount of pain; Patient satisfaction and analgesia; Patients will be asked 6, 12, and 24 hours after surgery, and the dose of general anesthesia will be measured and compared between the two groups.

Participants/Inclusion and exclusion criteria

Inclusion criteria: Patients over 18 years; Candidate for lumpectomy or partial mastectomy; Perform surgery under general anesthesia. Exclusion criteria: Prohibition of any nerve block; History of bleeding disease; Drug allergy; History of liver or kidney failure; Obesity (BMI above 35); Pregnancy or lactation; History of drugs addiction; History of neurological diseases; Infection of block; History of breast surgery; patients over 60 years.

Intervention groups

Intervention group 1: Type II pectoral nerve block.
Intervention group 2: Anterior Ceratus block under ultrasound guide.

Main outcome variables

The amount of pain; Patient satisfaction; Patient analgesia; Total dose of anesthesia

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20211215053419N1**

Registration date: **2022-02-26, 1400/12/07**

Registration timing: **prospective**

Last update: **2022-02-26, 1400/12/07**

Update count: **0**

Registration date

2022-02-26, 1400/12/07

Registrant information

Name

Parinaz Morovati sharif abadi

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 4608 8077

Email address

parinazmorovati43589@gmail.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2022-03-06, 1400/12/15

Expected recruitment end date

2022-07-23, 1401/05/01

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparison of efficacy of ultrasound guided pectoral nerve block (PECS II) versus serratus anterior block (SAB) on postoperative pain in breast cancer surgery

Public title

Efficacy of ultrasound guided pectoral nerve block versus serratus anterior block on pain

Purpose

Other

Inclusion/Exclusion criteria

Inclusion criteria:

Patients over 18 years old Candidate for lumpectomy or partial mastectomy Perform surgery under general anesthesia

Exclusion criteria:

Prohibition of any nerve block History of bleeding disease Drug allergy History of liver or kidney failure Obesity Pregnancy or lactation History of drugs addiction History of neurological diseases Infection of the site of the block History of breast surgery Patients over 60 years old

Age

From **18 years** old to **60 years** old

Gender

Female

Phase

N/A

Groups that have been masked

- Participant
- Data analyser

Sample size

Target sample size: **80**

Randomization (investigator's opinion)

Randomized

Randomization description

Patients will be randomly assigned to two groups by using a random number table. In order to randomize, a random block method will be used. For this purpose, we formed blocks of size 4. In each block, 2 individuals will be in intervention group 1 and 2 will be in intervention group 2. A total of 20 blocks will be considered for the study. Random Block Quadruple All possible blocks are arranged as follows: block 1: ABAB block 2: AABB block 3: ABBA block 4: BBAA block 5: BABA block 6: BAAB We need 20 blocks to select 80 people. We randomly select these blocks from 1 to 6. Using the software, we choose a random number between the numbers 1 to 6. For example, if the number 6 is selected as the first block and the number 2 as the second block, the people who enter the study will be given BAABAABB, respectively. Finally, group A receives control intervention and group B receives treatment intervention.

Blinding (investigator's opinion)

Single blinded

Blinding description

Patients will be unaware and blind to the type of treatment. Similar envelopes were used for hiding, without the name tag of the block method and only with the code. Patients will be aware that they will be randomly assigned to one of two treatment groups, but will not know which treatment will be offered in that

group. Patients will be placed in one of two groups using a table of random numbers. The data collector and analyst will collect and analyze the information based on groups 1 and 2, and will not be aware of the type of treatment provided in the groups and will be kept blind.

Placebo

Not used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Iran University of Medical Sciences

Street address

Iran University of Medical Sciences, Hemmat Highway

City

Tehran

Province

Tehran

Postal code

1449614535

Approval date

2021-12-04, 1400/09/13

Ethics committee reference number

IR.IUMS.FMD.REC.1400.528

Health conditions studied

1

Description of health condition studied

Breast cancer

ICD-10 code

C50

ICD-10 code description

Malignant neoplasm of breast

Primary outcomes

1

Description

The amount of pain

Timepoint

6, 12 and 24 hours after surgery

Method of measurement

Based on Numerical Rating Scale

2

Description

Patient satisfaction

Timepoint

6, 12 and 24 hours after surgery

Method of measurement

Based on Likert scale

3

Description

Patient analgesia

Timepoint

6, 12 and 24 hours after surgery

Method of measurement

Ramsay sedation scale

4

Description

Total dose of anesthesia

Timepoint

After surgery

Method of measurement

Based on patient records

Secondary outcomes

empty

Intervention groups

1

Description

Intervention group 1: Type II pectoral nerve block. The block will be performed after general anesthesia. The purpose of this block is to penetrate the two fascia compartments by dividing the dose of local anesthesia between the pectoral nerve in the pectoralis minor muscle. This block is performed by placing the patient supine and placing the surgical side arm in a 90 degree position. The linear transducer is located on the outer side of the midline of the clavicle at the level of the third rib, and in the first injection approximately 0.2 ml per kg of ropivacaine is injected 0.25 between the pectoralis major and minor muscles, for lateral and medial pectoral nerve blocks. The transducer is rotated slightly to create a path for the needle from the top and inside to the side. While the second injection was in the axillary anterior line and at the level of the fourth gear with 0.2 ml per kg of ropivacaine 0.25 between the pectoralis minor and the anterior ceratus muscle, using a linear transducer with a depth of 1 to 4 cm and a needle. Local block is performed. The injection depth is usually 1 to 3 cm for the first injection and 3 to 6 cm for the second injection.

Category

Treatment - Surgery

2

Description

Intervention group 2:Anterior Ceratus block under ultrasound guide. The block will be performed after general anesthesia under an ultrasound guide with a portable ultrasound device. In this block the linear

transducer is located on the midaxillary line and at the level of the fifth gear. A block needle is inserted from the outer edge of the transducer and moves forward until the tip of the needle reaches each plate between the thoracic fascia on the ultrasound image. Following the detection of ultrasound signs, a linear transducer with a depth of 1 to 4 cm and a local block needle can be performed and an injection of 0.4 ml per kg of ropivacaine 0.25 block will be performed. Device model: Portable Sono Site, HFL50X/15-6 MHz Trancducer, Made in USA.

Category

Treatment - Surgery

Recruitment centers

1

Recruitment center

Name of recruitment center

Hazrat Rasoul Akram hospital

Full name of responsible person

Parinaz Morovati Sharif Abadi

Street address

Hazrate Rasool hospital, Niyayesh street, Sattarkhan

City

Tehran

Province

Tehran

Postal code

1445613131

Phone

+98 21 6435 2190

Email

hrmc@iums.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Iran University of Medical Sciences

Full name of responsible person

Hossein Keyvani

Street address

Iran University of Medical Sciences, Hemmat Highway

City

Tehran

Province

Tehran

Postal code

1449614535

Phone

+98 21 8670 2503

Email

research@iums.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Iran University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding*empty***Country of origin****Type of organization providing the funding**

Academic

Street address

Hazrate Rasool hospital, Niyayesh street, Sattarkhan

City

Tehran

Province

Tehran

Postal code

1445613131

Phone

+98 21 4608 8077

Email

parinazmorovati43589@gmail.com

Person responsible for general inquiries**Contact****Name of organization / entity**

Iran University of Medical Sciences

Full name of responsible person

Parinaz Morovati Sharif Abadi

Position

resident

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Street address

Hazrate Rasool hospital, Niyayesh street, Sattarkhan

City

Tehran

Province

Tehran

Postal code

1445613131

Phone

+98 21 4608 8077

Email

parinazmorovati43589@gmail.com

Person responsible for updating data**Contact****Name of organization / entity**

Iran University of Medical Sciences

Full name of responsible person

Parinaz Morovati Sharif Abadi

Position

resident

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Street address

Hazrate Rasool hospital, Niyayesh street, Sattarkhan

City

Tehran

Province

Tehran

Postal code

1445613131

Phone

+98 21 4608 8077

Email

parinazmorovati43589@gmail.com

Person responsible for scientific inquiries**Contact****Name of organization / entity**

Iran University of Medical Sciences

Full name of responsible person

Parinaz Morovati Sharif Abadi

Position

resident

Latest degree

Medical doctor

Other areas of specialty/work

Anesthesiology

Sharing plan**Deidentified Individual Participant Data Set (IPD)***empty***Study Protocol***empty***Statistical Analysis Plan***empty***Informed Consent Form***empty***Clinical Study Report***empty***Analytic Code***empty***Data Dictionary***empty*