

Clinical Trial Protocol

Iranian Registry of Clinical Trials

02 Jul 2026

Evaluation of sweet aromatherapy effect on pain level of maxillary primary molars infiltration anesthesia in 7-9 years old children

Protocol summary

Study aim

evaluation the effect of sweet odor aromatherapy in pain level of maxillary molars infiltration in children 7-9 years old

Design

A clinical trial with a control group, with parallel groups, double-blind, randomized on 32 patients, a simple randomization method with a sealed envelope will be used for randomization.

Settings and conduct

Children aged 7-9 years in the pediatric ward of Qazvin Dental School who need maxillary molar infiltration included due to inclusion and exclusion criteria. Injection in the first intervention group is done in the room with saturated sweet smell, in the second intervention group, it is done at the same time with exposure to sweet smell (with a higher saturation than the first intervention group) and in the control group in an odorless environment. The evaluator encounters the child after preparation (evaluator blinding) and the statistical analyzer receives the data in group codes (analyzer blinding)

Participants/Inclusion and exclusion criteria

Children aged 7-9 years who need infiltration of maxillary primary molars, who are healthy and cooperative, are included in the study and are excluded from the study in case of any allergies or problems with sweets, smelling disorders or acute dental conditions.

Intervention groups

In the first session, the children receive brushing and fluoride therapy in order to shape their behavior, and in the next session, after using the tell-show-do technique, Injection in the first intervention group is done in the room with saturated sweet smell, in the second intervention group, it is done at the same time with exposure to sweet smell (with a higher saturation than the first group) and in the control group in an odorless environment.

Main outcome variables

Objective evaluation according to FLACC criteria;
Subjective evaluation according to Wong Baker criteria;
Heart rate; Degree of blood oxygen saturation

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20180104038218N3**

Registration date: **2022-04-09, 1401/01/20**

Registration timing: **prospective**

Last update: **2022-04-09, 1401/01/20**

Update count: **0**

Registration date

2022-04-09, 1401/01/20

Registrant information

Name

Razieh Jabbarian

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 28 3335 3061

Email address

r.jabbarian@qums.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2022-04-30, 1401/02/10

Expected recruitment end date

2022-07-01, 1401/04/10

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date
empty

Scientific title
Evaluation of sweet aromatherapy effect on pain level of maxillary primary molars infiltration anesthesia in 7-9 years old children

Public title
Evaluation of the effect of sweet odor on pain level of upper jaw anesthesia in children 7-9 years old

Purpose
Treatment

Inclusion/Exclusion criteria
Inclusion criteria:
children 7-9 years old requiring maxillary molar infiltration systemic and developmental health Positive behavior according to Frankle classification (cooperative and without resistance to treatment) Do not eat or drink anything for at least one hour before the injection
Exclusion criteria:
ear, nose, throat disorders allergy, sinusitis or tonsillitis current cold or any other smelling disorder The presence of spontaneous / nocturnal pain or abscess in the desired tooth Pain, inflammation and any other problems in the injection site child's previous dental experiment

Age
From **7 years** old to **9 years** old

Gender
Both

Phase
N/A

Groups that have been masked

- Outcome assessor
- Data analyser

Sample size
Target sample size: **48**

Randomization (investigator's opinion)
Randomized

Randomization description
Randomization will be done in blocked randomization method and blocking and sequencing of samples will be done with random allocation software; The names of the groups are then placed in envelopes from 1 to 48 and placed in a box, respectively. The researcher who will make the random assignment will not be aware of the contents of the envelopes and the contents of the envelopes indicate the study groups (anesthesia in the saturated environment of odor, sudden exposure to odor and control) and after ensuring that the sample enters the research and Obtaining written consent, the first envelope is removed in order and placed in one of the study groups according to the content of the envelope.

Blinding (investigator's opinion)
Double blinded

Blinding description
Outcome assessment is performed by a person other than the researcher and clinical caregiver and because he / she is present at the patient's bedside after the intervention, he / she is blind to the received

intervention. The grouping results of each sample are delivered to the statistical analyzer in the form of numbers.

Placebo
Not used

Assignment
Parallel

Other design features

Secondary Ids
empty

Ethics committees

1

Ethics committee
Name of ethics committee
Ethics Committee of Qazvin University of Medical Sciences
Street address
Bahonar Boulevard
City
Qazvin
Province
Qazvin
Postal code
3419759811

Approval date
2022-01-23, 1400/11/03

Ethics committee reference number
IR.QUMS.REC.1400.427

Health conditions studied

1

Description of health condition studied
injection in pediatric dentistry

ICD-10 code
G89.1

ICD-10 code description
Acute pain, not elsewhere classified

Primary outcomes

1

Description
objective level of pain

Timepoint
after completion of anesthesia

Method of measurement
Face,leg,activity,crying,consolability (FLACC)

2

Description
subjective pain level

Timepoint
after injection completion

Method of measurement

Wong baker

3

Description

heart rate

Timepoint

before, during and after injection

Method of measurement

pulse oximeter

4

Description

oxygen saturation

Timepoint

before, during and after injection

Method of measurement

pulse oximeter

Secondary outcomes

empty

Intervention groups

1

Description

Intervention group: In the first session, children receive brushing and fluoride therapy to shape their behavior, and in the next session, after using the tell-show-do technique, they are injected with infiltration in the area of the maxillary molars. The intervention group is placed in a room saturated with a sweet smell during injection. The injection technique in all three groups was the same, first by applying a topical benzocaine gel (20%) to the mucosa for one minute and then a carpol of 8.1 ml of 2% lidocaine + epinephrine 1: 80000 (Daroupakhsh, Iran) and a needle. 25 mm with gauge 27 (Sirio, Italy). After pulling the mucus, a needle is inserted at a 45 degree angle of 1-2 mm into the mucosa and a few drops are injected. After aspiration, more progress is made and the injection is completed. All injections are given by one person.

Category

Treatment - Other

2

Description

Intervention group: In the first session, children receive brushing and fluoride therapy to shape their behavior, and in the next session, after using the tell-show-do technique, they are injected with infiltration in the area of the maxillary molars. The second intervention group is exposed to a sudden sweet odor (with a higher saturation than the first intervention group) just during the injection. The injection technique in all three groups was the same, first by applying a topical benzocaine gel (20%) to the mucosa for one minute and then a carpol of 8.1 ml of 2% lidocaine + epinephrine 1: 80000

(Daroupakhsh, Iran) and a needle. 25 mm with gauge 27 (Sirio, Italy). After pulling the mucus, a needle is inserted at a 45 degree angle of 1-2 mm into the mucosa and a few drops are injected. After aspiration, more progress is made and the injection is completed. All injections are given by one person.

Category

Treatment - Other

3

Description

Control group: In the first session, children receive brushing and fluoride therapy to shape their behavior, and in the next session, after using the tell-show-do technique, they are injected with infiltration in the area of the maxillary molars. The control group is placed in the injection room and is not exposed to any odor. The injection technique in both groups was the same, first by applying a topical benzocaine gel (20%) to the mucosa for one minute and then a carpol of 8.1 ml of 2% lidocaine + epinephrine 1: 80000 (Daroupakhsh, Iran) and a needle. 25 mm with gauge 27 (Sirio, Italy). After pulling the mucus, a needle is inserted at a 45 degree angle of 1-2 mm into the mucosa and a few drops are injected. After aspiration, more progress is made and the injection is completed. All injections are given by one person.

Category

Treatment - Other

Recruitment centers

1

Recruitment center

Name of recruitment center

Qazvin university of medical sciences, faculty of dentistry

Full name of responsible person

Razieh Jabbarian

Street address

Bahonar Boulevard

City

Qazvin

Province

Qazvin

Postal code

Razieh Jabbarian

Phone

+98 28 3335 3062

Email

r.jabbarian@qums.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Qazvin University of Medical Sciences

Full name of responsible person

Dr Mehdi Mirhashemi

Street address

Shahid Beheshti boulevard

City

Qazvin

Province

Qazvin

Postal code

13911/34156

Phone

+98 28 7777 3333

Email

sm.mirhashemi@qums.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

No

Title of funding source

Qazvin university of medical sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Other

Person responsible for general inquiries

Contact

Name of organization / entity

Qazvin University of Medical Sciences

Full name of responsible person

Razieh Jabbarian

Position

assistant professor

Latest degree

Specialist

Other areas of specialty/work

Dentistry

Street address

Bahonar Boulevard

City

Qazvin

Province

Qazvin

Postal code

3471976160

Phone

+98 28 3335 3062

Email

r.jabbarian@qums.ac.ir

Person responsible for scientific inquiries

Contact

Name of organization / entity

Qazvin University of Medical Sciences

Full name of responsible person

Razieh Jabbarian

Position

assistant professor

Latest degree

Specialist

Other areas of specialty/work

Dentistry

Street address

Bahonar Boulevard

City

Qazvin

Province

Qazvin

Postal code

3471976160

Phone

+98 28 3335 3062

Email

r.jabbarian@qums.ac.ir

Person responsible for updating data

Contact

Name of organization / entity

Qazvin University of Medical Sciences

Full name of responsible person

Razieh Jabbarian

Position

assistant professor

Latest degree

Specialist

Other areas of specialty/work

Dentistry

Street address

Bahonar Boulevard

City

Qazvin

Province

Qazvin

Postal code

3471976160

Phone

+98 28 3335 3062

Email

r.jabbarian@qums.ac.ir

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to

make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available