

Clinical Trial Protocol

Iranian Registry of Clinical Trials

04 Jun 2026

Evaluation of adding Lidocaine to Morphine in patient controlled analgesia on pain intensity after unilateral knee prosthesis surgery by spinal anesthesia

Protocol summary

Study aim

We measure pain reduction after unilateral knee arthroplasty after adding lidocaine to morphine in a patient controlled analgesia pump.

Design

Clinical trial with a control group, with parallel groups, three blinded, randomized, phase 2-3 on 58 patients.

Settings and conduct

This study was conducted in Aria and Farabi hospitals in Mashhad. A pain pump was placed for the patients for post-operative analgesia. The pain level of the surgical site was compared with the Visual Analogue Scale method at specific time intervals. In this study, drugs were used with the same packaging and blinding was done for patients, project personnel and researchers.

Participants/Inclusion and exclusion criteria

Inclusion criteria: Patients with American Society of Anesthesiologists (ASA) class 1 and 2 between 45 and 75 years old, with a Body Mass Index(BMI) in the range of 20-25 should undergo spinal anesthesia for the surgery:
Exclusion criteria: Patients with a history of epilepsy, known kidney disease, heart block, or drug addiction, Patients with a medical history that indicates hypersensitivity to lidocaine and opioids, Patients with coagulation disorders, Patients with skin infection at the injection site, a history Moderate to severe nausea and vomiting in the past 24 hours

Intervention groups

Group A: The content of the patient controlled analgesia pump attached to the patient included 20 mg of morphine, 8 mg of Ondansetron ampoule (4 cc), 1g of Apotel (6.7 cc) and the rest of normal saline to reach a total volume of 100 cc
Group B: The content of the patient controlled analgesia pump includes 10 mg of morphine, 50 cc of lidocaine 2%, 8 mg of ampoule of Ondansetron (4 cc), 1g of Apotel (6.7 cc) and the rest of normal saline to a total volume of 100 cc.

Main outcome variables

Postoperative pain; mean arterial pressure; heart rate; nausea and vomiting; systolic pressure; diastolic pressure

General information

Reason for update

Acronym

PCA

IRCT registration information

IRCT registration number: **IRCT20200718048125N2**

Registration date: **2022-12-14, 1401/09/23**

Registration timing: **registered_while_recruiting**

Last update: **2022-12-14, 1401/09/23**

Update count: **0**

Registration date

2022-12-14, 1401/09/23

Registrant information

Name

Hamed Beyzaii

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 51 3765 5378

Email address

hmd_beyzaii@yahoo.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2022-11-27, 1401/09/06

Expected recruitment end date

2023-03-19, 1401/12/28

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Evaluation of adding Lidocaine to Morphine in patient controlled analgesia on pain intensity after unilateral knee prosthesis surgery by spinal anesthesia

Public title

The effect of adding lidocaine to morphine on pain intensity after unilateral knee prosthesis surgery

Purpose

Treatment

Inclusion/Exclusion criteria**Inclusion criteria:**

Patients who are in American Society of Anesthesiologists(ASA) class 1 and 2 Patients who are between 45 and 75 years old Patients with a Body Mass Index(BMI) in the range of 20-25 Patients who undergo spinal anesthesia for surgery

Exclusion criteria:

Patients with a history of epilepsy Patients with known kidney disease Patients with heart block Patients with drug addiction Patients with medical histories that show sensitivity to lidocaine and opioid substances Patients with coagulation disorders Patients with skin infection at the injection site Patients who had a history of moderate to severe nausea and vomiting in the last 24 hours

Age

From **45 years** old to **75 years** old

Gender

Both

Phase

2-3

Groups that have been masked

- Participant
- Care provider
- Investigator
- Outcome assessor
- Data analyser
- Data and Safety Monitoring Board

Sample size

Target sample size: **58**

Randomization (investigator's opinion)

Randomized

Randomization description

A simple randomization method is used. We assign an A and B code to each anesthetic intervention. With the lottery method, patients will be placed in one of two codes. In this way, 29 codes A and 29 codes B are written on the paper and each paper is placed inside a special non-transparent sealed envelope and the envelopes are randomly placed on top of each other. Upon the entry of each qualified patient, an envelope is returned by the anesthesiologist to that patient and according to the code written on the paper inside, the anesthetic method is applied.

Blinding (investigator's opinion)

Triple blinded

Blinding description

In this study, patients are placed in two separate groups, A and B, which have the same drug composition except for one type of drug, and drugs with the same shape and packaging are injected to the patients through a patient controlled analgesia pump. The patient does not know whether he is in the control group or the intervention group. The personnel cooperating in the plan do not know the content of the pain pump. The researcher also has no knowledge of which group each patient is in and only analyzes the information.

Placebo

Not used

Assignment

Parallel

Other design features**Secondary Ids**

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Ethics committee of Islamic Azad University, Mashhad branch

Street address

No. 8, Emam Khomeini Ave, Shahinfar Medical School

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Approval date

2022-11-16, 1401/08/25

Ethics committee reference number

IR.IAU.MSHD.REC.1401.157

Health conditions studied**1****Description of health condition studied**

Osteoarthritis of knee

ICD-10 code

M17.9

ICD-10 code description

Osteoarthritis of knee, unspecified

Primary outcomes**1****Description**

Postoperative pain

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Visual Analogue Scale

2

Description

Heart rate

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Pulse oximetry

3

Description

Mean arterial pressure

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Clinical examination

4

Description

Postoperative nausea and vomiting

Timepoint

4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Clinical examination

5

Description

Oxygen saturation

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Pulse oximetry

6

Description

Systolic pressure

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Dial sphygmomanometer

7

Description

Diastolic pressure

Timepoint

Immediately after the surgery and 4 hours after the surgery, 8 hours after the surgery, 24 hours after the surgery

Method of measurement

Dial sphygmomanometer

Secondary outcomes

empty

Intervention groups

1

Description

Control group: Spinal anesthesia; 25-gauge spinal needle (Dr.J Tianjin Hanaco Medical Co. Tianjin, China) was introduced, then 0.5% Bupivacaine mixed with 0.2 mg of Adrenaline was injected at either the third or fourth lumbar epidural space with median or paramedian approach. The content of the patient controlled analgesia pump attached to the patient included 20 mg of morphine, 8 mg of Ondansetron ampoule (4 cc), 1g of Apotel (6.7 cc) and the rest of normal saline to reach a total volume of 100 cc.

Category

Treatment - Drugs

2

Description

Intervention group: Spinal anesthesia; 25-gauge spinal needle (Dr.J Tianjin Hanaco Medical Co. Tianjin, China) was introduced, then 0.5% Bupivacaine mixed with 0.2 mg of Adrenaline was injected at either the third or fourth lumbar epidural space with median or paramedian approach. The contents of the patient controlled analgesia pump include 10 mg of morphine, 50 cc of 2.8% lidocaine, 8 mg of Ondansetron ampoule (4 cc), one gram of Apotel (6.7 cc) and the rest of normal saline to reach a total volume of 100 cc.

Category

Treatment - Drugs

Recruitment centers

1

Recruitment center

Name of recruitment center

Farabi hospital

Full name of responsible person

Hamed Beyzaee

Street address

No.1, Kowsar Blvd, Mashhad.

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3807 2000

Fax

+98 51 3881 7980

Email

hmd_beyzaii@yahoo.com

2

Recruitment center

Name of recruitment center

Aria hospital

Full name of responsible person

Hamed Beyzaee

Street address

No. 3, Pasdaran Ave, Chamran street, Mashhad.

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3222 9094

Email

hmd_beyzaii@yahoo.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Islamic Azad University

Full name of responsible person

Tooraj Zandbaf

Street address

Shahinfar Faculty of Medicine, Imam Khomeini Ave

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3225 6291

Email

IMRC@mshdiau.ac.ir

Web page address

<http://med.mshdiau.ac.ir/>

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Islamic Azad University

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Islamic Azad University

Full name of responsible person

Hamed Beyzaee

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Shahinfar faculty of medicine, Imam Khomeini Ave

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3225 6291

Email

hmd_beyzaii@yahoo.com

Web page address

<http://med.mshdiau.ac.ir/>

Person responsible for scientific inquiries

Contact

Name of organization / entity

Islamic Azad University

Full name of responsible person

Hamed Beyzaee

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

No 48. 6th baharestan st, Sadjad blv.

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3765 5378

Email

hmd_beyzaii@yahoo.com

Person responsible for updating data

Contact

Name of organization / entity

Islamic Azad University

Full name of responsible person

Hamed Beyzaee

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

No 48. 6th baharestan st, Sadjad blv.

City

Mashhad

Province

Razavi Khorasan

Postal code

9187147578

Phone

+98 51 3765 5378

Email

hmd_beyzaii@yahoo.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

No - There is not a plan to make this available

Justification/reason for indecision/not sharing IPD

There is no further information.

Study Protocol

No - There is not a plan to make this available

Statistical Analysis Plan

No - There is not a plan to make this available

Informed Consent Form

No - There is not a plan to make this available

Clinical Study Report

No - There is not a plan to make this available

Analytic Code

No - There is not a plan to make this available

Data Dictionary

No - There is not a plan to make this available