

Clinical Trial Protocol

Iranian Registry of Clinical Trials

10 Jun 2026

Clinical and biomechanical comparison of microscopic bilateral laminotomy versus total laminectomy with posterior spinal fusion for decompression of central canal and lateral recess simultaneous stenosis, with a focus on multilevel stenosis: A prospective comparative study.

Protocol summary

Study aim

To compare the clinical and biomechanical outcomes of microscopic bilateral laminotomy versus total laminectomy+ PSF in treating multi-level lumbar spinal stenosis (LSS)

Design

A two-arm parallel randomized interventional trial without blinding, with a sample size of 76 patients.

Settings and conduct

The present study will conduct on 76 patients in Shohada Tajrish Hospital. Patients will be randomized into intervention groups: A (Microscopic bilateral laminotomy) and B (Total laminectomy+ PSF).

Participants/Inclusion and exclusion criteria

Inclusion criteria: patients between 30-75 with ≥ 3 levels LSS; stenosis of the central canal (mild to moderate), and lateral recess simultaneously, who suffers from resistant radicular pain as the primary complaint. Exclusion criteria: patients with severe stenosis of the central canal who suffer from severe neurogenic claudication that limits their ability; extruded discs or those requiring discectomy; osteoporotic patients T score < -1.5 ; previous history of laminotomy, laminectomy, or PSF 5. Serious medical conditions that preclude surgical procedures; current metabolic, rheumatologic, or inflammatory disorders requiring corticosteroid consumption; spinal deformity.

Intervention groups

Intervention group 1: patients with multi-level spinal stenosis, simultaneously stenosis of the central canal and lateral recess, who will be scheduled to undergo microscopic bilateral laminotomy. Intervention group 2: patients with multi-level spinal stenosis, simultaneously stenosis of the central canal and lateral recess, who will be scheduled to undergo total laminectomy with PSF (bilaterally removal of the lamina ≥ 3 , spinous processes,

interspinous intertransverse ligaments and partial medial facetectomy, as well as PSF)

Main outcome variables

Changes in Visual Analogue Scale and Pelvic incidence-lumbar lordosis mismatch

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20190126042496N3**

Registration date: **2023-02-01, 1401/11/12**

Registration timing: **registered_while_recruiting**

Last update: **2023-02-01, 1401/11/12**

Update count: **0**

Registration date

2023-02-01, 1401/11/12

Registrant information

Name

Mohammadrez Shahmohammadi

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 2271 8001

Email address

mr_shahmohammadi@sbmu.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2023-01-25, 1401/11/05

Expected recruitment end date

2023-07-27, 1402/05/05

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Clinical and biomechanical comparison of microscopic bilateral laminotomy versus total laminectomy with posterior spinal fusion for decompression of central canal and lateral recess simultaneous stenosis, with a focus on multilevel stenosis: A prospective comparative study.

Public title

Clinical and biomechanical outcomes following decompression of central canal and lateral recess simultaneous stenosis, with a focus on multilevel stenosis: A randomized comparison of microscopic bilateral laminotomy versus total laminectomy with posterior spinal fusion.

Purpose

Treatment

Inclusion/Exclusion criteria**Inclusion criteria:**

Between 30- and 75-year-old age Multi-level canal stenosis (≥ 3 levels) Stenosis of both the central canal (mild to moderate) and the lateral recess simultaneously. Failure of conservative therapy for radicular pain after six months. Radicular pain predominates over claudication(mild).

Exclusion criteria:

Patients with severe stenosis of the central canal. Patients with severe neurogenic claudication suffer from leg pain that limits their ability to stand or walk. Extruded discs or those requiring discectomy Osteoporotic patients with a T score < -1.5 Previous spinal surgery history (laminotomy , laminectomy, or posterior spinal fusion) Serious medical conditions such as congestive heart failure, cirrhosis, and other chronic diseases that preclude surgical procedures. Even though patients develop cardiac, renal, or pulmonary disorders during the study, they will be excluded. Current metabolic or rheumatologic and inflammatory disorders requiring corticosteroid consumption. Spinal deformity

AgeFrom **30 years** old to **75 years** old**Gender**

Both

Phase

N/A

Groups that have been masked*No information***Sample size**Target sample size: **76****Randomization (investigator's opinion)**

Randomized

Randomization description

By using a random number table (simple randomization), all patients who meet all the inclusion and exclusion

criteria will be consecutively recruited from outpatient clinics and randomized into one of the study arms by the statistician. (1:1)

Blinding (investigator's opinion)

Not blinded

Blinding description**Placebo**

Not used

Assignment

Parallel

Other design features**Secondary Ids**

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Institutional Research Ethics Committee - Shahid Beheshti University of Medical Sciences

Street address

3rth floor, Faculty of Medicine, next to Taleghani Hospital, Evin, Shahid Chamran Highway

City

Tehran

Province

Tehran

Postal code

1985717434 - 1939547

Approval date

2021-01-26, 1399/11/07

Ethics committee reference number

IR.SBMU.MSP.REC.1399.655

Health conditions studied**1****Description of health condition studied**

Patients with multilevel spinal canal stenosis, simultaneous stenosis of the central canal, and lateral recess.

ICD-10 code

M48.0

ICD-10 code description

Spinal stenosis

Primary outcomes**1****Description**

VAS changes : Post-op versus pre-op VAS changes for leg pain and back pain

Timepoint

VAS scores will be measured before surgery, after 6-9 months, and after 18-24 months to determine the extent of pain relief.

Method of measurement

The patient's pain level is assessed on a horizontal line of 100 mm that its left end represents no pain, and the right end corresponds to the most severe pain that can be experienced. The patient is asked to mark the pain intensity they are experiencing currently.

2**Description**

Duration of hospitalization

Timepoint

Total number of hospital admission days

Method of measurement

Number of days staying in hospital with reference to medical records

3**Description**

Duration of operation

Timepoint

During surgery

Method of measurement

The number of hours it takes for the operation to be completed.

4**Description**

Blood transfusion: The number of patients requiring blood transfusions.

Timepoint

During surgery

Method of measurement

Based on doctor's diagnosis.

5**Description**

Wound infection and discharge: Incidence of postoperative wound infection and discharge

Timepoint

Immediately after surgery until the end of the hospitalization period for each patient (up to 1 month after surgery)

Method of measurement

Based on doctor's diagnosis.

6**Description**

Incidental durotomy: An accidental puncture or laceration of dura during a procedure.

Timepoint

During surgery

Method of measurement

Based on doctor's diagnosis.

7**Description**

meningitis: Incidence of meningitis after surgery

Timepoint

Immediately after surgery until the end of the hospitalization period for each patient (up to 1 month after surgery)

Method of measurement

Based on doctor's diagnosis.

8**Description**

Revision surgery: The number of patients requiring revision surgery either due to technical issues or infections.

Timepoint

Immediately after surgery until the end of the follow-up time

Method of measurement

Based on doctor's diagnosis.

9**Description**

Root injury: Incidence of root injury during surgery

Timepoint

During surgery

Method of measurement

Based on doctor's diagnosis.

10**Description**

Vacuum sign: Incidence of a collection of gas in the intervertebral disc space

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

Based on doctor's diagnosis via lateral plain radiographs.

11**Description**

Adjacent segment disease (ASD): Degeneration of the mobile spinal segments between upper instrumented vertebra (UIV) and UIV+2 leading to spinal canal stenosis or discopathy .

Timepoint

After 18-24 months follow-up

Method of measurement

Anteroposterior and lateral plain radiographs, and computed tomography (CT)

12**Description**

Hypermobility: Increased range of motion between UIV and UIV+2

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

Flexion-extension radiographs.

Secondary outcomes

1

Description

Proximal junctional kyphosis (PJK): A rise of over 20° in the Cobbs' angle between the lower endplate of the upper instrumented vertebra (UIV) and the upper endplates of two super-adjacent vertebrae between the immediate postoperative and after follow-up time

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral radiograph

2

Description

Sagittal vertical axis (SVA): The length of a horizontal line connecting the superior-posterior endplate of S1 to a vertical plumb line that drawn from the mid-point of the C7 vertebral body. This horizontal distance should be located within \pm 5mm of the superior-posterior endplate of S1.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral spine radiographs

3

Description

Pelvic incidence (PI): The angle is formed by the perpendicular line running from the midpoint of the sacral plate and a line connecting that point to the center of the bicoxofemoral axis that serves as the foundation for all other spinal curves.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral radiograph

4

Description

Pelvic tilt (PT): The angle between the vertical axis and the line connecting the center of the coxofemoral axis to the midpoint of the S1 endplate.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral radiograph

5

Description

Sacral slope (SS): The angle between the horizontal reference line and the line parallel to the superior plate of S1.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral radiograph

6

Description

lumbar lordosis (LL): The angle between the caudal endplate of L5 and the upper plane of the L1 lumbar vertebra.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The standing lateral radiograph

7

Description

Pelvic incidence-lumbar lordosis mismatch (DPILL = PI-LL): A mismatch between pelvic incidence and lumbar lordosis strongly correlates with the sagittal balance and has become a key element in treating adult deformities.

Timepoint

Immediately after surgery, after 6-9 months, and after 18-24 months follow-up

Method of measurement

The difference between pelvic incidence and lumbar lordosis

Intervention groups

1

Description

Intervention group 1 (Group A): A group of patients with multi-level spinal canal stenosis, simultaneously stenosis of the central canal and lateral recess, who will be scheduled to undergo lumbar decompression surgery via microscopic bilateral laminotomy through a unilateral approach.

Category

Treatment - Surgery

2

Description

Intervention group 2 (Group B): A group of patients with multi-level spinal canal stenosis, simultaneously stenosis of the central canal and lateral recess, who will be scheduled to undergo lumbar decompression surgery via total laminectomy and partial medial facetectomy with posterior spinal fusion (bilaterally removal of the bony lamina \geq 3 lamina, spinous processes, and interspinous intertransverse ligaments and medial facet, as well as posterior spinal fusion with pedicle screw fixation to prevent instability)

Category

Treatment - Surgery

Recruitment centers

1

Recruitment center

Name of recruitment center

Shohada-e-Tajrish hospital

Full name of responsible person

Mohammadreza Shahmohammadi

Street address

Shohada Tajrish Hospital, Tajrish Square, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1989934148

Phone

+98 21 25719

Email

mr_shahmohammadi@sbmu.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Neurofunctional Research Center of Shohada Tajrish Hospital

Full name of responsible person

Prof. Dr. Afshin Zarghi

Street address

Neurofunctional Research Center of Shohada Tajrish Hospital, Tajrish Square, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

19839-63113

Phone

+98 21 2270 1022

Email

neurofunctional.cntr@gmail.com

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Neurofunctional Research Center of Shohada Tajrish Hospital

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Shahid Beheshti University of Medical Sciences

Full name of responsible person

Mohammadreza Shahmohammadi

Position

Assistant Professor

Latest degree

Specialist

Other areas of specialty/work

Neurosurgery

Street address

Next to Taleghani hospital, Shahid Araabi St., Yaman St., Shahid Chamran highway, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1985717443

Phone

+98 21 23871

Email

mr_shahmohammadi@sbmu.ac.ir

Person responsible for scientific inquiries

Contact

Name of organization / entity

Shahid Beheshti University of Medical Sciences

Full name of responsible person

Mohammadreza Shahmohammadi

Position

Assistant Professor

Latest degree

Specialist

Other areas of specialty/work

Neurosurgery

Street address

Next to Taleghani hospital, Shahid Araabi St., Yaman St., Shahid Chamran highway, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1985717443

Phone

+98 21 23871

Email

mr_shahmohammadi@sbmu.ac.ir

Person responsible for updating data

Contact

Name of organization / entity

Shahid Beheshti University of Medical Sciences

Full name of responsible person

Mohammadreza Shahmohammadi

Position

Assistant Professor

Latest degree

Specialist

Other areas of specialty/work

Neurosurgery

Street address

Next to Taleghani hospital, Shahid Araabi St., Yaman St., Shahid Chamran highway, Tehran, Iran

City

Tehran

Province

Tehran

Postal code

1985717443

Phone

+98 21 23871

Email

mr_shahmohammadi@sbmu.ac.ir

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available