

Clinical Trial Protocol

Iranian Registry of Clinical Trials

23 Jul 2026

Investigating the preemptive effect of two different doses of dexmedetomidine on changes in blood pressure and heart rate following the use of a tourniquet during a randomized trial of upper extremity orthopedic surgery

Protocol summary

Study aim

Determining the effect of two different doses of 1 and 0.75 ($\mu\text{g/kg}$) dexmedetomidine on the control of heart rate and blood pressure during the use of tourniquet in upper extremity surgery

Design

Clinical trial with control group, parallel groups, triple-blind, randomized, phase 3 on 90 patients
Randomization with random allocation software

Settings and conduct

In Kashani hospital in Isfahan, under EKG monitoring, non-invasive blood pressure and pulse oximetry Dex group 1, 1 ($\mu\text{g/kg}$) dexmedetomidine induced within 10 minutes, continuation with 0.75 ($\mu\text{g/kg}$) Dex group 2, 1 ($\mu\text{g/kg}$) dexmedetomidine induced within 10 minutes, continuation with 1 ($\mu\text{g/kg}$) Saline group (control), normal saline infusion Identical syringes are used Induction of anesthesia using sodium thiopental 5 (mg/kg), fentanyl 3 ($\mu\text{g/kg}$), atracurium 0.6 (mg/kg) Maintenance of anesthesia using propofol 0.75 (min/ $\mu\text{g/kg}$) and remifentanyl 0.1 (min/ $\mu\text{g/kg}$) Measurement and recording of systolic, diastolic, and mean arterial blood pressure, heart rate, and SpO₂ at baseline and at 0, 10, 20, 30, 40, 50, and 60 minutes after inflating the tourniquet and right after emptying it The patient, the person who collects the data, and the statistician are blinded.

Participants/Inclusion and exclusion criteria

Inclusion criteria: patients with ASA (American Society of Anesthesiologists) = 1, aged 18-65 years Exclusion criteria: drug addicts, people with a history of allergy to studied drugs, people with heart, liver and kidney failure, allergy to study drugs, change in anesthesia method, duration of tourniquet less than 60 or more than 150 minutes

Intervention groups

Three groups of 30 patients, dexmedetomidine 1 and 2

and the control group Dex group 1: dose 0.75 ($\mu\text{g/kg}$) of dexmedetomidine infusion Dex group 2: dose 1 ($\mu\text{g/kg}$) of dexmedetomidine infusion Control group: normal saline infusion

Main outcome variables

Heart rate; Blood pressure

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20090129001615N7**

Registration date: **2023-05-13, 1402/02/23**

Registration timing: **prospective**

Last update: **2023-05-13, 1402/02/23**

Update count: **0**

Registration date

2023-05-13, 1402/02/23

Registrant information

Name

Azim Honarmand

Name of organization / entity

Alzahra hospital

Country

Iran (Islamic Republic of)

Phone

+98 31 3668 0048

Email address

honarmand@med.mui.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2023-05-21, 1402/02/31

Expected recruitment end date

2023-11-21, 1402/08/30

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Investigating the preemptive effect of two different doses of dexmedetomidine on changes in blood pressure and heart rate following the use of a tourniquet during a randomized trial of upper extremity orthopedic surgery

Public title

Investigating the preemptive effect of two different doses of dexmedetomidine on changes in blood pressure and heart rate following the use of a tourniquet during a randomized trial of upper extremity orthopedic surgery

Purpose

Prevention

Inclusion/Exclusion criteria**Inclusion criteria:**

Patients with ASA (American Society of Anesthesiologists) = 1, aged 18-65 years, who are Patients aged 18-65 years Candidates for upper limb orthopedic surgery under tourniquet who undergo general anesthesia

Exclusion criteria:

Addicted people People with a history of allergy to the studied drugs People with heart, liver or kidney failure People with ASA = 2

Age

From **18 years** old to **65 years** old

Gender

Both

Phase

3

Groups that have been masked

- Participant
- Outcome assessor
- Data analyser

Sample size

Target sample size: **90**

Randomization (investigator's opinion)

Randomized

Randomization description

It will be done using Random Allocation software. Patients are placed in three groups of 30, dexmedetomidine 1 and 2 and saline (control) group.

Blinding (investigator's opinion)

Triple blinded

Blinding description

The patient and the person collecting the study data are not aware of the study group. The statistician who analyzes the data will also include this requirement.

Placebo

Used

Assignment

Parallel

Other design features**Secondary Ids**

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Ethics Committee of Isfahan University of Medical Sciences

Street address

Soffeh st.

City

Isfahan

Province

Isfahan

Postal code

8174675731

Approval date

2023-03-14, 1401/12/23

Ethics committee reference number

IR.ARI.MUI.REC.1401.334

Health conditions studied**1****Description of health condition studied**

Blood pressure

ICD-10 code**ICD-10 code description****2****Description of health condition studied**

Heart rate

ICD-10 code**ICD-10 code description****Primary outcomes****1****Description**

Blood pressure

Timepoint

At baseline times (before administration of study drugs) and then at times 0, 10, 20, 30, 40, 50 and 60 minutes after tourniquet inflation and then immediately after deflation

Method of measurement

Sphygmomanometer

2**Description**

Heart rate
Timepoint
At baseline times (before administration of study drugs) and then at times 0, 10, 20, 30, 40, 50 and 60 minutes after tourniquet inflation and then immediately after deflation
Method of measurement
Electrocardiogram

3

Description
Oxygen saturation

Timepoint
At baseline times (before administration of study drugs) and then at times 0, 10, 20, 30, 40, 50 and 60 minutes after tourniquet inflation and then immediately after deflation
Method of measurement
Pulse oximeter

Secondary outcomes

empty

Intervention groups

1

Description
Intervention group one: In the dexmedetomidine 1 group, 1 (kg/μg) dexmedetomidine is infused within 10 minutes and then continued with a dose of 0.75 (kg/μg) dexmedetomidine infusion during the operation.

Category
Prevention

2

Description
Intervention group two: In the dexmedetomidine group 2, 1 (kg/μg) dexmedetomidine is infused within 10 minutes and then continued with a dose of 1 (kg/μg) dexmedetomidine infusion during the operation.

Category
Prevention

3

Description
Control group: In the control group, normal saline is infused.

Category
Placebo

Recruitment centers

1

Recruitment center
Name of recruitment center
Ayatollah Kashani Hospital
Full name of responsible person

Azim Honarmand
Street address
Kashani st.
City
Isfahan
Province
Isfahan
Postal code
8174673461
Phone
+98 31 3233 0091
Email
honarmand@med.mui.ac.ir

Sponsors / Funding sources

1

Sponsor
Name of organization / entity
Esfahan University of Medical Sciences
Full name of responsible person
Gholamreza Askari
Street address
Hezar Jerib Ave.
City
Isfahan
Province
Isfahan
Postal code
8174673461
Phone
+98 31 3668 0048
Email
gh_askari@med.mui.ac.ir
Grant name
Grant code / Reference number
Is the source of funding the same sponsor organization/entity?
Yes
Title of funding source
Esfahan University of Medical Sciences
Proportion provided by this source
100
Public or private sector
Public
Domestic or foreign origin
Domestic
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
Academic

Person responsible for general inquiries

Contact
Name of organization / entity
Esfahan University of Medical Sciences
Full name of responsible person
Azim Honarmand
Position

Anesthesiologist and special care specialist

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Hezar Jerib Ave.

City

Isfahan

Province

Isfahan

Postal code

8174673461

Phone

+98 913 110 2327

Fax

Email

honarmand@med.mui.ac.ir

Web page address

Person responsible for scientific inquiries

Contact

Name of organization / entity

Esfahan University of Medical Sciences

Full name of responsible person

Azim Honarmand

Position

Anesthesiologist and special care specialist

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Hezar Jerib Ave.

City

Isfahan

Province

Isfahan

Postal code

8174673461

Phone

+98 913 110 2327

Fax

Email

honarmand@med.mui.ac.ir

Web page address

Person responsible for updating data

Contact

Name of organization / entity

Esfahan University of Medical Sciences

Full name of responsible person

Azim Honarmand

Position

Anesthesiologist and special care specialist

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Hezar Jerib Ave.

City

Isfahan

Province

Isfahan

Postal code

8174673461

Phone

+98 913 110 2327

Fax

Email

honarmand@med.mui.ac.ir

Web page address

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Undecided - It is not yet known if there will be a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available