

Clinical Trial Protocol

Iranian Registry of Clinical Trials

21 Jun 2026

Comparison of ovarian response to ovulation induction after laparoscopic cystectomy or medical management among infertile women with endometrioma undergoing assisted reproductive technology

Protocol summary

Summary

The purpose of this study is to compare the ovarian response to ovulation induction after laparoscopic cystectomy or medical management among infertile women suffering from endometrioma going through assisted reproductive technology. Our main objective is to define the exact number of oocytes retrieved after these two processes. The sample includes all infertile women aged 40 years or less, with asymptomatic endometrioma undergoing assisted reproductive technology. Diagnosis of endometrioma is based on the evidence found through trans vaginal ultrasound indicating round shaped homogeneous hypoechoe tissue. Infertile couples with maternal cause who undergo TESE (Testicular Sperm Extraction) or PESA (Percutaneous Epididimal Sperm Aspiration) are excluded from the current study. The sample includes 80 patients divided in two groups. First group involves patients with asymptomatic endometrioma receiving IM Dipherelin prior to ovulation induction and the second group are the patients undergoing ovulation induction after laparoscopic ovarian cystectomy followed by IM Dipherelin injection. All patients in both groups receive a single IM dose of Dipherelin monthly for 3 consecutive months before the start of ovulation induction by Gonadotropins. After observing at least two 18 mm follicles in trans vaginal ultrasound, 10 000 units, hCG (10000 units, IM) is injected. The following step is ovarian puncture guided by trans vaginal ultrasound under general anesthesia 36 hours later. Trans cervical embryo transfer is performed 3 days after the puncture. In this study, the following outcomes are compared between two groups: number of retrieved oocytes, dominant follicles, embryos; quality of embryos; the rate of fertilization, chemical and clinical pregnancy.

General information

Acronym

IRCT registration information

IRCT registration number: **IRCT201106116689N2**
Registration date: **2012-11-05, 1391/08/15**
Registration timing: **registered_while_recruiting**

Last update:

Update count: **0**

Registration date

2012-11-05, 1391/08/15

Registrant information

Name

Sedigheh Hosseinimousa

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 8800 8810

Email address

hoseinimosa@sina.tums.ac.ir

Recruitment status

Recruitment complete

Funding source

Vice Chancellor for Research, Tehran University of Medical Sciences

Expected recruitment start date

2012-01-04, 1390/10/14

Expected recruitment end date

2012-12-30, 1391/10/10

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparison of ovarian response to ovulation induction after laparoscopic cystectomy or medical management among infertile women with endometrioma undergoing assisted reproductive technology

Public title

Treatment of endometrioma in infertile women

Purpose

Treatment

Inclusion/Exclusion criteria

Inclusion criteria: Infertile women with endometrioma requiring assisted reproductive technology Exclusion criteria: Age over 40 years; Endometrioma smaller than 2 cm or larger than 6 cm; Male infertility TESE (Testicular Sperm Extraction) or PESA (Percutaneous Epididimal Sperm Aspiration)

Age

From **20 years** old to **40 years** old

Gender

Female

Phase

N/A

Groups that have been masked

No information

Sample size

Target sample size: **80**

Randomization (investigator's opinion)

Randomized

Randomization description

Blinding (investigator's opinion)

Not blinded

Blinding description

Placebo

Not used

Assignment

Parallel

Other design features

All volunteers with inclusion criteria are asked to participate in the current study if agreed, a informed consent form is signed. she is randomly assigned to one of two groups based in Bertoli distribution. If she has an objection, we will switch her to another group.

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethic Committee of Tehran University of Medical Sciences

Street address

Ethic Committee, Tehran University of Medical Sciences, Poorsina Street, Tehran

City

Tehran

Postal code

Approval date

2012-01-02, 1390/10/12

Ethics committee reference number

90-02-30-14261-40996

Health conditions studied

1

Description of health condition studied

Infertility

ICD-10 code

N97

ICD-10 code description

ناباروري زنانه

2

Description of health condition studied

Endometriomas

ICD-10 code

N80.1

ICD-10 code description

اندومتریوز بزرگ تخمدان

Primary outcomes

1

Description

The number of retrieved oocytes

Timepoint

84 days after the first Dipherelin injection

Method of measurement

Invert Microscope, Diaphot 300, Nikon, Japan

Secondary outcomes

1

Description

The number of dominant follicles

Timepoint

82 days after the first Dipherelin injection

Method of measurement

nGS= number of observed gestational sac in trans vaginal sonography, nET= number of embryonal transfer, clinical pregnancy rate(%)= nGS/ nET x100

2

Description

fertilization rate

Timepoint

87 days after the first Dipherelin injection

Method of measurement

nE= number of total embryos, nI= number of injected oocytes, fertilization rate(%)= nE/ nI x 100

3

Description

The number and quality of embryos

Timepoint

87 days after the first Dipherelin injection

Method of measurement

Invert Microscope, Diaphot 300, Nikon, Japan

4

Description

chemical pregnancy

Timepoint

101 days after the first Dipherelin injection

Method of measurement

βhCG test

5

Description

clinical pregnancy rate

Timepoint

115 days after the first Dipherelin injection

Method of measurement

nGS= number of observed gestational sac in trans vaginal sonography, nET= number of embryonal transfer, clinical pregnancy rate(%)= nGS/ nET x100

Intervention groups

1

Description

First group receives 3 consecutive single IM doses of Dipherelin 3.75mg (Beaufouipfen, France) in 3 consecutive months. Ten days after the third IM dose of Dipherelin ovarian stimulation is conducted by using Gonal F (Serono, Switzerland, 300-450IU daily), for one week. Then Gonal F is replaced by HMG (Ferring, Germany, 300-450IU daily) until the observation of 18mm follicles in trans vaginal ultrasound. For visualizing follicular development, trans vaginal ultrasound (Sonoline G20; Siemens Medical Solutions, California, USA) is performed every 4 days. After the observation of at least two 18mm follicles, HCG (Ferring Co, Germany, 10000IU, IM) is injected and after 36 hours oocyte retrieve is performed under general anesthesia. Three days after fertilization technique performed through intra cytoplasmic sperm injection (ICSI), trans cervical embryo transfer will be carried out. Detection of pregnancy is through serum BhCG analysis, 3 days after embryonic transfer and the clinical pregnancy is detected by the aid of trans vaginal ultrasound, two weeks later when the pregnancy sac is detected. Main outcome is the number of oocytes retrieved. The number of oocytes retrieved, dominant follicles, embryos; quality of embryos; the rate of fertilization, laboratory and clinical pregnancy are compared between two groups.

Category

Treatment - Drugs

2

Description

Second group receives 3 consecutive single IM doses of Dipherelin 3.75mg (Beaufouipfen, France) in 3 consecutive months after laparoscopic ovarian cystectomy. Ten days after the third IM dose of Dipherelin ovarian stimulation is conducted by using Gonal F (Serono, Switzerland, 300-450IU daily), for one week. Then Gonal F is replaced by HMG (Ferring, Germany, 300-450IU daily) until the observation of 18mm follicles in trans vaginal ultrasound. For visualizing follicular development, trans vaginal ultrasound (Sonoline G20, Siemens Medical Solutions, California, USA) is performed every 4 days. After observing at least two 18mm follicles, HCG (Ferring Co, Germany, 10000IU, IM) is injected and after 36 hours oocyte retrieve is performed under general anesthesia. Three days after fertilization technique performed through intra cytoplasmic sperm injection (ICSI), trans cervical embryo transfer is carried out. Detection of pregnancy is through serum BhCG analysis, 3 days after embryonic transfer and the clinical pregnancy is detected by the aid of trans vaginal ultrasound, two weeks later when the pregnancy sac is detected. Main outcome is the number of oocytes retrieved. The number of oocytes retrieved, dominant follicles, embryos; quality of embryos; the rate of fertilization, laboratory and clinical pregnancy are compared between two groups.

Category

Treatment - Surgery

Recruitment centers

1

Recruitment center

Name of recruitment center

Infertility Unit, Shariati Hospital

Full name of responsible person

Dr Sedighe Hoseinimosa

Street address

Department of Infertility, Shariati Hospital, Tehran University of Medical Sciences, North Karegar Street

City

Tehran

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Vice Chancellor for Research, Tehran University of Medical Sciences

Full name of responsible person

Dr Akbar Fotouhi

Street address

Vice Chancellor for Research, Tehran University of Medical Sciences, Keshavarz Boulevard, Tehran

City

Tehran

Grant name
Grant code / Reference number
Is the source of funding the same sponsor organization/entity?
Yes
Title of funding source
Vice Chancellor for Research, Tehran University of Medical Sciences
Proportion provided by this source
100
Public or private sector
empty
Domestic or foreign origin
empty
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
empty

Person responsible for general inquiries

Contact

Name of organization / entity
Tehran University of Medical Sciences, Shariati Hospital, Department of Infertility
Full name of responsible person
Dr Sedighe Hoseinimosa
Position
Infertility Fellowship
Other areas of specialty/work
Street address
Infertility Department, Shariati Hospital, North Karegar Street, Tehran
City
Tehran
Postal code
1411713135
Phone
+98 21 8800 8810
Fax
+98 21 8822 0050
Email
hoseinimosa@razi.tums.ac.ir
Web page address

Person responsible for scientific inquiries

Contact

Name of organization / entity
Tehran University of Medical Sciences
Full name of responsible person
Dr Leili Safdarian
Position
Associated Professor

Other areas of specialty/work
Street address
Infertility Department, Shariati Hospital, North Karegar Street, Tehran
City
Tehran
Postal code
1411713135
Phone
+98 21 8800 8810
Fax
+98 21 8822 0050
Email
safdarian@sina.tums.ac.ir
Web page address

Person responsible for updating data

Contact

Name of organization / entity
Tehran University of Medical Sciences
Full name of responsible person
Dr Sedighe Hoseinimosa
Position
Infertility Fellowship
Other areas of specialty/work
Street address
Infertility Unit, Shariati Hospital, North Karegar Street, Tehran
City
Tehran
Postal code
1411713135
Phone
+98 21 8800 8810
Fax
00982188220050
Email
hoseinimosa@razi.tums.ac.ir
Web page address

Sharing plan

Deidentified Individual Participant Data Set (IPD)
empty
Study Protocol
empty
Statistical Analysis Plan
empty
Informed Consent Form
empty
Clinical Study Report
empty
Analytic Code
empty
Data Dictionary
empty