

Clinical Trial Protocol

Iranian Registry of Clinical Trials

10 Jun 2026

Comparing the effects of bupivacaine and bupivacaine-methylprednisolone in ultrasound guided erector spinae plane block on postoperative pain in Lumbar spine surgery

Protocol summary

Study aim

Comparing the effects of bupivacaine and bupivacaine-methylprednisolone in ultrasound guided erector spinae plane block on post operative pain in Lumbar spine surgery

Design

Clinical trial with control and intervention group, single blind, on 64 patients, randomized with sealed envelope.

Settings and conduct

Patients referred to Luqman Hospital are divided into two intervention and control groups of 32 people by block randomization. After anesthesia with the same method, both groups will be placed in the prone position before the surgery under ultrasound guidance under bilateral erector spina block at the level of the surgical site. Paramedian sagittal ultrasound probe, about 2 cm outside the spinous processes, we find the transverse process on the same side. We insert the needle caudal to the cranial so that the tip of the needle hits the transverse process. If the needle site is suitable, 20 cc of bupivacaine-methylprednisolone will be injected in the intervention group and 20 cc of bupivacaine 0.25% in the control group. Isoflurane and opioid consumption during surgery, pain, nausea and vomiting during recovery and sugar measurement The blood will be determined up to 24 hours later, as well as the pain level of the patient up to one month.

Participants/Inclusion and exclusion criteria

Inclusion criteria : Patients 18-65 years old; Normal kidney and liver function; ASA score 1-2; Patient consent ; No history of allergies to local anesthetics; No drug addiction Exclusion criteria:, Increase the scope of surgery to more than three level ,time of surgery for more than 6 hours; No diabetes

Intervention groups

The intervention group, after anesthesia and changing to the prone position, ,are subjected to erector spinae block

with bupivacaine-methylprednisolone, and the control group are subjected to block with bupivacaine

Main outcome variables

Consumption of isofluran , opioids, pain score

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20210415050983N6**

Registration date: **2024-04-18, 1403/01/30**

Registration timing: **prospective**

Last update: **2024-04-18, 1403/01/30**

Update count: **0**

Registration date

2024-04-18, 1403/01/30

Registrant information

Name

Sogol Asgari

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 8836 3185

Email address

drasgari98429@sbmu.ac.ir

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2024-05-21, 1403/03/01

Expected recruitment end date

2024-08-22, 1403/06/01

Actual recruitment start date

empty

Actual recruitment end date
empty

Trial completion date
empty

Scientific title
Comparing the effects of bupivacaine and bupivacaine-methylprednisolone in ultrasound guided erector spinae plane block on postoperative pain in Lumbar spine surgery

Public title
Comparing the effects of bupivacaine and bupivacaine-methylprednisolone in ultrasound guided erector spinae plane block on postoperative pain in Lumbar spine surgery

Purpose
Other

Inclusion/Exclusion criteria
Inclusion criteria:
Patients 18-65 years old are candidates for two or three level spine surgery ASA score 1-2 Normal kidney and liver function Patient consent to perform the block No history of allergies to local anesthetics No drug addiction No diabetes
Exclusion criteria:
Increase the scope of surgery to more than three level Extending the length of surgery for more than 6 hours Block site infection or systemic History of anticoagulant use

Age
From **18 years** old to **65 years** old

Gender
Both

Phase
3

Groups that have been masked

- Participant
- Care provider
- Outcome assessor
- Data analyser

Sample size
Target sample size: **64**

Randomization (investigator's opinion)
Randomized

Randomization description
Permuted Randomized Blocks :In this method, 10 random blocks are generated by computer. Each block includes 5 people in the intervention group and 5 people in the control group. The order of these people is randomly arranged by computer and people are assigned to groups in the same way. At the end of each block, a new block of 10 is produced and this process will continue until the final sample volume is reached.

Blinding (investigator's opinion)
Single blinded

Blinding description
Participants in the study are unaware of the groupings because the intervention is performed after anesthesia. The patient's clinical caregiver, the evaluator and recorder of the results, and the data analyzer are not aware of the grouping.

Placebo
Not used

Assignment
Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Vice for Research and Technology, Shahid Beheshti University of Medical Sciences

Street address

Velenjak, Yemen Street, Shahid Shahriari Square

City

Tehran

Province

Tehran

Postal code

1985717443

Approval date

2024-01-24, 1402/11/04

Ethics committee reference number

IR.SBMU.MSP.REC.1402.575

Health conditions studied

1

Description of health condition studied

Lumbar discopathy

ICD-10 code

M51.36

ICD-10 code description

Other intervertebral disc degeneration, lumbar region

Primary outcomes

1

Description

Intraoperative isoflurane Consumption in tow groups

Timepoint

Before the start of anesthesia and after the end of anesthesia

Method of measurement

By a graduated glass based on mL

2

Description

Intraoperative fentanyl consumption in the two groups

Timepoint

End of surgery

Method of measurement

Dosage consumed based on mcg

3

Description

Pain after surgery

Timepoint

0, 1 and 6 hours after surgery and one month after surgery

Method of measurement

NRS

4

Description

blood sugar

Timepoint

After surgery up to 24 months

Method of measurement

Blood test

Secondary outcomes

1

Description

nausea and vomiting

Timepoint

0, 1 and 6 hours after surgery

Method of measurement

Ask the patient

Intervention groups

1

Description

Control group: After anesthesia and change of position to peron before surgery, in sterile conditions, using peripheral nerve block needle (stimuQuik, ARROW use) with sonosite-Nerve Ultrasound system under the erector spina block One-sided and each-sided injection of 20 ml of 0.25% bupivacaine was performed by a trained anesthesiologist in accordance with standard guidelines. A 5-8MHz liner probe is used for nerve block ultrasound guides. After selecting the target process transducer, the sagittal paramedic prop is placed about 2 cm outside the spinous processes so that the process transducer can be seen in the same direction. Insert the needle inplean the codal from the cranial to cudal until the tip of the needle hits the process transducer. 1-2 cc of local anesthetic is injected to ensure the correct location of the needle. If the location of the needle is suitable, the medicine is injected. The needle point is towards the posterior and inferior side.

Category

Treatment - Drugs

2

Description

Intervention group: Intervention group: After anesthesia and change of position to peron before surgery, in sterile conditions, using peripheral nerve block needle (stimuQuik, ARROW use) with sonosite-Nerve Ultrasound

system under the erector spina block One-sided and each-sided injection of 20 ml of Bupivacaine and methylprednisolone 40 was performed by a trained anesthesiologist in accordance with standard guidelines. A 5-8MHz liner probe is used for nerve block ultrasound guides. After selecting the target process transducer, the sagittal paramedic prop is placed about 2 cm outside the spinous processes so that the process transducer can be seen in the same direction. Insert the needle inplean the codal from the cranial to cudal until the tip of the needle hits the process transducer. 1-2 cc of local anesthetic is injected to ensure the correct location of the needle. If the location of the needle is suitable, the medicine is injected. The needle point is towards the posterior and inferior side.

Category

Treatment - Drugs

Recruitment centers

1

Recruitment center

Name of recruitment center

Loghman Hakim Hospital

Full name of responsible person

Sogol Asgari

Street address

South Kargar St. - Kamali St. - Special St.

City

Tehran

Province

Tehran

Postal code

1333635445

Phone

+98 21 5102 5291

Email

drasgari98429@gmail.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Shahid Beheshti University of Medical Sciences

Full name of responsible person

Dr. Seyed Ali Ziaei

Street address

Velenjak, Yemen Street, Shahid Shahriari Square

City

Tehran

Province

Tehran

Postal code

1985717443

Phone

+98 21 23871

Email

aliziai@sbmu.ac.ir

Grant name

Grant code / Reference number
Is the source of funding the same sponsor organization/entity?
No
Title of funding source
Shahid Beheshti University of Medical Sciences
Proportion provided by this source
100
Public or private sector
Public
Domestic or foreign origin
Domestic
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
Academic

Person responsible for general inquiries

Contact
Name of organization / entity
Shahid Beheshti University of Medical Sciences
Full name of responsible person
Sogol Asgari
Position
Assistant Professor
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
South Kargar St. Kamali St. Special Loghman Hakim Hospital
City
Tehran
Province
Tehran
Postal code
1333635445
Phone
+98 21 5541 9005
Email
Drasgari98429@gmail.com

Person responsible for scientific inquiries

Contact
Name of organization / entity
Shahid Beheshti University of Medical Sciences
Full name of responsible person
Sogol Asgari
Position
Assistant Professor
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
South Kargar St. Kamali St. Special Loghman Hakim

Hospital
City
Tehran
Province
Tehran
Postal code
1333635445
Phone
+98 21 5541 9005
Email
Drasgari98429@gmail.com

Person responsible for updating data

Contact
Name of organization / entity
Shahid Beheshti University of Medical Sciences
Full name of responsible person
Sogol Asgari
Position
Assistant Professor
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
Velenjak, Yemen Street, Shahid Shahriari Square
City
Tehran
Province
Tehran
Postal code
1985717443
Phone
+98 21 23871
Email
Drasgari98429@gmail.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)
Undecided - It is not yet known if there will be a plan to make this available
Study Protocol
Undecided - It is not yet known if there will be a plan to make this available
Statistical Analysis Plan
Undecided - It is not yet known if there will be a plan to make this available
Informed Consent Form
Undecided - It is not yet known if there will be a plan to make this available
Clinical Study Report
Undecided - It is not yet known if there will be a plan to make this available
Analytic Code
Undecided - It is not yet known if there will be a plan to make this available
Data Dictionary
Undecided - It is not yet known if there will be a plan to make this available