

Clinical Trial Protocol

Iranian Registry of Clinical Trials

01 Jun 2026

Comparison of outcomes between Propofol and Dexmedetomidine on extubation time and vital signs in patients with post- coronary artery bypass grafting in Open heart Intensive Care Unit

Protocol summary

Study aim

Overall Aim is to compare the effect of Dexmedetomidine and Propofol on extubation time and vital signs in patients undergoing coronary artery bypass grafting in the Intensive Care Unit Open Heart. The practical goal of this project is to select the right medication for sedation of post-CABG patients with fewer side effects, more efficient and more suitable for extubation.

Design

Clinical trial with a total of two parallel and equal groups, double-blind, randomized, on 92 patients. A block method was used for randomization

Settings and conduct

CABG candidate patients who are referred to Amirul Mominin Arak Hospital. Double-blind collection of vital signs and other variables. The patients are divided into two groups by anesthesiologist so the patient and the evaluator are not aware of the drug's name used in the groups

Participants/Inclusion and exclusion criteria

Terms and conditions of participation: Having informed consent Patients must meet inclusion criteria, i.e. be elective and referred for open heart surgery only. Age should be between 35 and 80 years old. Performed by the same surgeon. The maximum duration of the surgery must be 6 hours. All were operated with a cardiopulmonary pump. Valve surgery should not be performed at the same time as CABG Exclusion conditions are: Patient is not a candidate for elective CABG surgery. Duration of surgery is more than 6 hours. Patients have not had surgery outside of the cardiopulmonary pump. Patients outside the 35-80 age range. The patient has the right to withdraw at any stage of the research

Intervention groups

Two equal groups of 46 people. One group receives

propofol medication and the other group receives dexmedetomidine medication, then the effect of these two medications on these two groups is measured.

Main outcome variables

Mean Arterial Pressure Pulse rate SpO2 Extubation time Hospitalization time Department hospitalization time

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20240207060924N1**

Registration date: **2024-02-17, 1402/11/28**

Registration timing: **prospective**

Last update: **2024-02-17, 1402/11/28**

Update count: **0**

Registration date

2024-02-17, 1402/11/28

Registrant information

Name

Taha Rafiei

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 86 3313 6055

Email address

taharafiei@yahoo.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2024-02-20, 1402/12/01

Expected recruitment end date

2024-11-21, 1403/09/01

Actual recruitment start date
empty

Actual recruitment end date
empty

Trial completion date
empty

Scientific title
Comparison of outcomes between Propofol and Dexmedetomidine on extubation time and vital signs in patients with post- coronary artery bypass grafting in Open heart Intensive Care Unit

Public title
Effect of Propofol in coronary bypass surgery

Purpose
Treatment

Inclusion/Exclusion criteria
Inclusion criteria:
Having informed consent Patients must meet inclusion criteria, i.e. be elective and referred for open heart surgery only Age should be between 35 and 80 years old Performed by the same surgeon The maximum duration of the surgery must be 6 hours All were operated with a cardiopulmonary pump Valve surgery should not be performed at the same time as CABG
Exclusion criteria:
Patient is not a candidate for elective CABG surgery Duration of surgery is more than 6 hours Patients have not had surgery outside of the cardiopulmonary pump Patients outside the 35-80 age range The patient has the right to withdraw at any stage of the research

Age
From **35 years** old to **80 years** old

Gender
Both

Phase
3

Groups that have been masked

- Participant
- Investigator

Sample size
Target sample size: **92**

Randomization (investigator's opinion)
Randomized

Randomization description
In this study, we will randomly divide 92 patients into two equal groups (P) and (D) using random block method. The randomization unit will be done individually and we will use statistical software as a tool. All statistical combinations that these two groups can have will be written. These combinations will include PPDD, PDDP, PDPD, DDPP, DPDP, DPPD. Now, using these combinations, the observations are placed into two intervention groups. Note that the intervention groups are chosen blindly.

Blinding (investigator's opinion)
Double blinded

Blinding description
The aforementioned study is a double-blind study in

which patients who meet the inclusion criteria and provide informed consent are assigned to one of two groups receiving Propofol (P) or Dexmedetomidine (D). Patients are randomized into groups (P) and (D) by the anesthesiologist and are unaware of which group they are in. In addition, the intern in charge of the project does not know the names of the groups. After the patients wake up and become conscious and are extubated while the sedative infusion is stopped, the student in charge of the project will be present in the ward to fill in the questionnaires, so he will not know about the studied groups and will be blinded.

Placebo
Not used

Assignment
Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Arak University of Medical Sciences

Street address

Basij square, sardasht, Arak university of medical science

City

Arak

Province

Markazi

Postal code

3819693345

Approval date

2024-02-07, 1402/11/18

Ethics committee reference number

IR.ARAKMU.REC.1402.212

Health conditions studied

1

Description of health condition studied

Comparison of Outcomes Between Propofol and Dexmedetomidine in Post-Coronary Artery Bypass Graft Patients

ICD-10 code

ICD-10 code description

Primary outcomes

1

Description

Mean arterial blood pressure

Timepoint

Measurement of mean arterial blood pressure at 5, 15, 30, 45, 60, 90, 120, 150 minutes after entering the ward

Method of measurement

Pulse Oximeter

2

Description

Pulse rate

Timepoint

Measurement of pulse rate at 5, 15, 30, 45, 60, 90, 120, 150 minutes after entering the ward

Method of measurement

Pulse Oximeter

3

Description

Oxygen saturation

Timepoint

Measurement of Oxygen saturation at 5, 15, 30, 45, 60, 90, 120, 150 minutes after entering the ward

Method of measurement

Pulse Oximeter

4

Description

Extubation time

Timepoint

The duration of extubation from the time of admission to the Open Heart Intensive Care Unit

Method of measurement

Questionnaire

5

Description

Length of stay in the Unit

Timepoint

The duration of the stay in the Intensive Care Unit of the Open Heart surgery after the operation

Method of measurement

Questionnaire

6

Description

Length of hospitalization

Timepoint

The duration of the stay in the hospital

Method of measurement

Questionnaire

7

Description

Ramsay Score

Timepoint

Measurement of Ramsay score at 5, 15, 30, 45, 60, 90, 120, 150 minutes after admission to the unit

Method of measurement

Questionnaire

8

Description

Mortality

Timepoint

Measurement of the number of mortality events in the study population

Method of measurement

Questionnaire

9

Description

Age

Timepoint

Registration of patient age

Method of measurement

Questionnaire

10

Description

Sex

Timepoint

Registration of patient sex

Method of measurement

Questionnaire

Secondary outcomes

empty

Intervention groups

1

Description

First intervention group: (P) that we infuse 10 cc per hour (10 cc/hr) for 4 to 6 hours from a vial of propofol 10% (manufactured by B. Braun, Germany).

Category

Treatment - Drugs

2

Description

Second intervention group: (D) In the Dexmedetomidine group, the syringe pump is filled to a volume of 50 cc and Dexmedetomidine is infused at a dose of 20 µg/hour (manufactured by Elixir Lorestan, Iran) for 4 to 6 hours.

Category

Treatment - Drugs

Recruitment centers

1

Recruitment center

Name of recruitment center

Amir Al Mo'menin Hospital

Full name of responsible person

Alireza Kamali

Street address

Amir Al Mo'menin Hospital, University campus, Basij square, Sardasht, Markazi, Iran

City

Arak

Province

Markazi

Postal code

3848176941

Phone

+98 86 3836 0000

Fax

+98 86 3417 3619

Email

it-amiralmomenin@arakmu.ac.ir

Position

Student of medicine

Latest degree

A Level or less

Other areas of specialty/work

General Practitioner

Street address

Faculty of medicine, Basij square, Sardasht

City

Arak

Province

Markazi

Postal code

3819693345

Phone

+98 86 3313 3147

Email

taharafiei@yahoo.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Arak University of Medical Sciences

Full name of responsible person

Alireza Kamali

Street address

Faculty of Medicine, Basij square, Sardasht

City

Arak

Province

Markazi

Postal code

3819693345

Phone

+98 86 3313 6055

Email

info@arakmu.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Arak University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Arak University of Medical Sciences

Full name of responsible person

Taha Rafiei

Person responsible for scientific inquiries

Contact

Name of organization / entity

Arak University of Medical Sciences

Full name of responsible person

Alireza Kamali

Position

Associate Professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Faculty of medicine, Basij square, Sardasht

City

Arak

Province

Markazi

Postal code

3819693345

Phone

+98 86 3313 6055

Email

alikalaliir@yahoo.com

Person responsible for updating data

Contact

Name of organization / entity

Arak University of Medical Sciences

Full name of responsible person

Taha Rafiei

Position

Student of medicine

Latest degree

A Level or less

Other areas of specialty/work

General Practitioner

Street address

Faculty of medicine, Basij square, Sardasht

City

Arak
Province
Markazi
Postal code
3819693345
Phone
+98 86 3313 6055
Email
info@arakmu.ac.ir

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to make this available

Analytic Code

Yes - There is a plan to make this available

Data Dictionary

Not applicable

Title and more details about the data/document

Data will be shared after study completion in accordance with ethical considerations

When the data will become available and for how long

Upon completion of the study

To whom data/document is available

The public is allowed access

Under which criteria data/document could be used

For scientific, research and therapeutic purposes

From where data/document is obtainable

first author

What processes are involved for a request to access data/document

By mentioning the reference of the study

Comments