

Clinical Trial Protocol

Iranian Registry of Clinical Trials

27 Jul 2026

Study of Preemptive Analgesic effect of dexmedetomidine Versus ketamine on pain control after total laparoscopic hysterectomy - a randomized clinical trial study

Protocol summary

Study aim

To evaluate and compare demographic data, post-operative pain intensity, opioid consumption, side effects, quality of recovery, hospital stay duration, intraoperative vital signs, and extubation time among the three study groups. (ketamine, dexmedetomidine and control)

Design

A randomized, double-blind, parallel-group, controlled phase III clinical trial was conducted on 120 patients. Randomization was performed using the card-shuffling method.

Settings and conduct

A randomized, double-blind clinical trial was conducted on 120 TLH patients at Hajar Hospital to compare dexmedetomidine and ketamine against a control. Drugs were administered 10 minutes before incision. Outcomes—including VAS pain scores, opioid use, vital signs, recovery quality, and hospital stay—were monitored for 24 hours postoperatively. Blinding was maintained for patients, anesthesiologists, and statisticians, with only the preparation nurse remaining unblinded.

Participants/Inclusion and exclusion criteria

A: Inclusion Criteria: Patients scheduled for Total Laparoscopic Hysterectomy (TLH). Patients with ASA physical status I or II. No history of addiction to benzodiazepines or opioids, including methadone. B: Exclusion Criteria: Prolonged surgical duration exceeding 180 minutes. Conversion of the surgical procedure to laparotomy. Requirement for blood transfusion during surgery.

Intervention groups

In one group, ketamine at a dose of 0.5 mg/kg and in another group, dexmedetomidine at a dose of 0.5 microgram/kg are administered intravenously after induction of anesthesia and before the start of surgery.

In the control group, an equivalent volume of normal saline is administered

Main outcome variables

Postoperative pain intensity was assessed using the Visual Analog Scale (VAS) at 0, 30 minutes, 1 hour, and 4, 6, 12, and 24 hours post-surgery.

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20190127042511N3**

Registration date: **2026-06-10, 1405/03/20**

Registration timing: **prospective**

Last update: **2026-06-10, 1405/03/20**

Update count: **0**

Registration date

2026-06-10, 1405/03/20

Registrant information

Name

Alireza Babaeizadeh

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 31 3669 4561

Email address

alirezabbz@yahoo.com

Recruitment status

recruiting

Funding source

Expected recruitment start date

2026-06-22, 1405/04/01

Expected recruitment end date

2027-03-20, 1405/12/29

Actual recruitment start date
empty

Actual recruitment end date
empty

Trial completion date
empty

Scientific title
Study of Preemptive Analgesic effect of dexmedetomidine Versus ketamine on pain control after total laparoscopic hysterectomy - a randomized clinical trial study

Public title
A comparative study of the effects of ketamine and dexmedetomidine on pain after hysterectomy

Purpose
Prevention

Inclusion/Exclusion criteria
Inclusion criteria:
 Patients undergoing Total Laparoscopic Hysterectomy (TLH). Patients with an American Society of Anesthesiologists (ASA) physical status classification of I or II. No history of addiction to benzodiazepines or opioids (including methadone).
Exclusion criteria:
 Prolonged surgery duration (exceeding 180 minutes). Conversion from laparoscopic to open surgery (laparotomy). Requirement for blood transfusion during surgery.

Age
No age limit

Gender
Female

Phase
3

Groups that have been masked

- Participant
- Care provider
- Investigator
- Outcome assessor
- Data analyser

Sample size
Target sample size: **120**

Randomization (investigator's opinion)
Randomized

Randomization description
Sampling was performed by convenience sampling. Patients who met the inclusion criteria were enrolled in the study and then randomly assigned to three groups: K, D, and control. For randomization, the card-shuffling method was used. In this method, a number of cards was selected by the researcher for the first group, and the same number of cards was considered for the subsequent groups. The cards were then mixed together (shuffled), one card was drawn, and its allocation was recorded. After being drawn, the card was returned to the set of cards. The cards were then reshuffled, and another card was drawn. This process continued until a random sequence corresponding to the sample size was

obtained. One of the advantages of this method is its practicality for multigroup studies. Olson et al. also used this method for randomization in their study.

Blinding (investigator's opinion)
Double blinded

Blinding description
To maintain the double-blind nature of the study, a coding method was used so that neither the performing physician nor the patient was aware of the type of intervention. The medications were prepared and administered by a trained nurse who was not involved in any other part of the study, and the coding procedure was also performed by this nurse. The blinding process was carried out as follows: 1. Individuals blinded to the intervention Patients: were unaware of the type of intervention administered (active drug or placebo). The physician performing the procedure: was blinded to the type of intervention and the composition of the medication contained in the syringes. The statistical analyst: did not have access to the group allocation codes until completion of the primary analysis. 2. Individuals aware of the intervention type Only one trained nurse, who had no role in treatment administration, sampling, patient monitoring, or outcome assessment, was aware of the group allocations. This individual was solely responsible for preparing the medications and coding the syringes and had no involvement in any other stage of the study. 3. Preparation and coding procedure The non-involved nurse prepared the active drug or placebo according to the randomization list. Each syringe was labeled with a unique numerical code without any identifying marks. The volume, color, and appearance of the drug and placebo were made identical to prevent recognition by either the patient or the physician. The correspondence table between the code and intervention type was kept in a sealed envelope and was opened only after completion of data collection. 4. Injection procedure The physician received only the coded syringe and had no information about its contents. The patient also received no information regarding the type of intervention, and the appearance of all syringes was completely identical.

Placebo
Used

Assignment
Parallel

Other design features

Secondary Ids
empty

Ethics committees

1

Ethics committee
Name of ethics committee
 Research Ethics Committees of School of Medicine - Shahrekord University of Medical Sciences
Street address
 Parastar St., Hajar Hospital

City
Shahrekord
Province
Chahar-Mahal-va-Bakhtiari
Postal code
۸۸۱۶۷۵۴۶۳۳
Approval date
2026-04-22, 1405/02/02
Ethics committee reference number
IR.SKUMS.MED.REC.1405.049

Health conditions studied

1

Description of health condition studied

Study of Preemitive Analgesic effect of dexmedetomidine Versus ketamine on pain control after total laparoscopic hysterectomy

ICD-10 code

ICD-10 code description

Primary outcomes

1

Description

Postoperative pain intensity, measured via the Visual Analog Scale (VAS), was evaluated at 0, 30 minutes, and 1 hour post-surgery, and at 4, 6, 12, and 24 hours after transfer to the ward, across the dexmedetomidine, ketamine, and control groups

Timepoint

at 0, 30 minutes, and 1 hour postoperatively, and at 4, 6, 12, and 24 hours following ward transfer in the dexmedetomidine, ketamine, and control groups.

Method of measurement

Visual Analog Scalescores were assessed on a scale of 0-10 at the indicated time points.

Secondary outcomes

1

Description

Total opioid consumption

Timepoint

Total fentanyl or pethidine consumption over 24 hours

Method of measurement

miligram

2

Description

Postoperative adverse effects

Timepoint

Postoperative adverse effects, including nausea, vomiting, hypotension, and bradycardia, were monitored in all groups.

Method of measurement

Based on the yes/no responses

3

Description

Length of hospital stay

Timepoint

Duration of stay until discharge after surgery

Method of measurement

Time to discharge, measured in days or hours

4

Description

vital sign

Timepoint

Every 15 minutes during the procedure

Method of measurement

Using a monitoring device (blood oxygen saturation, blood pressure in mmHg, and heart rate in beats per minute).

5

Description

Quality of recovery

Timepoint

At the time of discharge from recovery.

Method of measurement

Aldrete score

6

Description

Extubation time

Timepoint

Time from discontinuation of the anesthetic agent to extubation.

Method of measurement

Time from discontinuation of the anesthetic agent to extubation, measured in minutes.

Intervention groups

1

Description

In the first intervention group, general anesthesia is induced using midazolam (0.1 mg/kg), propofol (2.5 mg/kg), fentanyl (3

Category

Prevention

2

Description

"In the second intervention group, general anesthesia was induced using midazolam (0.1 mg/kg), propofol (2.5 mg/kg), fentanyl (3

Category

Prevention

3

Description

In the control group, general anesthesia was induced using midazolam (0.1 mg/kg), propofol (2.5 mg/kg), fentanyl (3

Category

Placebo

Recruitment centers

1

Recruitment center

Name of recruitment center

Hajar hospital

Full name of responsible person

Alireza Babaeizadeh

Street address

Parastar St., Hajar Hospital

City

Shahre-kird

Province

Chahar-Mahal-va-Bakhtiari

Postal code

۸۸۱۶۷۵۴۶۳۳

Phone

+98 38 3222 0016

Email

Hajar-Hospital@skums.ac.ir

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Shahre-kord University of Medical Sciences

Full name of responsible person

Alireza babaeizadeh

Street address

parastar

City

shahrekord

Province

Chahar-Mahal-va-Bakhtiari

Postal code

8815713471

Phone

+98 38 3333 3815

Email

mail@mail.skums.ac.ir

Web page address

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

No

Title of funding source

Reseach department

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Academic

Person responsible for general inquiries

Contact

Name of organization / entity

Shahre-kord University of Medical Sciences

Full name of responsible person

Alireza Babaeizadeh

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Parastar St., Hajar Hospital

City

Shahrekord

Province

Chahar-Mahal-va-Bakhtiari

Postal code

۸۸۱۶۷۵۴۶۳۳

Phone

+98 38 2223 0016

Email

babaeizadeh.a@skums.ac.ir

Person responsible for scientific inquiries

Contact

Name of organization / entity

Shahre-kord University of Medical Sciences

Full name of responsible person

Alireza Babaeizadeh

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Parastar St., Hajar Hospital

City

Shahrekord

Province

Chahar-Mahal-va-Bakhtiari

Postal code

۸۸۱۶۷۵۴۶۳۳

Phone

+98 38 2223 0016

Email

babaeizadeh.a@skums.ac.ir

Person responsible for updating data

Contact

Name of organization / entity

Shahre-kord University of Medical Sciences

Full name of responsible person

Alireza Babaeizadeh

Position

Assistant professor

Latest degree

Specialist

Other areas of specialty/work

Anesthesiology

Street address

Parastar St., Hajar Hospital

City

Shahrekord

Province

Chahar-Mahal-va-Bakhtiari

Postal code

۸۸۱۶۷۵۴۶۳۳

Phone

+98 38 2223 0016

Email

babaeizadeh.a@skums.ac.ir

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Yes - There is a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Yes - There is a plan to make this available

Analytic Code

Yes - There is a plan to make this available

Data Dictionary

Yes - There is a plan to make this available

Title and more details about the data/document

De-identified raw dataset containing the variables used for statistical analyses in the present study. All direct patient identifiers will be removed prior to data sharing.

When the data will become available and for how long

From the date of publication until 5 years after publication.

To whom data/document is available

Qualified researchers who provide a scientifically justified request and agree to data confidentiality requirements.

Under which criteria data/document could be used

Data will be available for research and educational purposes only. Commercial use is prohibited. Recipients must agree not to attempt re-identification of study participants.

From where data/document is obtainable

Requests should be directed to the corresponding author via email and will be reviewed on a case-by-case basis.

What processes are involved for a request to access data/document

Requests for access to the data should be submitted in writing to the corresponding author. Requests will be reviewed by the research team for scientific merit, consistency with the study objectives, and compliance with ethical requirements. Upon approval, applicants will be required to sign a Data Use Agreement, after which de-identified data will be made available.

Comments